

THURSDAY, SEPTEMBER 22, 1898.

THE FAUNA AND FLORA OF THE PAMIR.

Report on the Natural History Results of the Pamir Boundary Commission. By A. W. Alcock, M.B., Surgeon Naturalist to the Commission. With a List of the Plants, by J. F. Duthie, B.A., F.L.S., and a Notice of the Rock-specimens, by T. H. Holland, A.R.C.S., F.G.S. (Calcutta, 1898.)

FOR this useful addition to our knowledge of the Pamir plateau we are indebted to the enlightened policy of the Government of India, who, in attaching Mr. Alcock as naturalist to the Commission despatched in 1895 to demarcate the boundary between Afghan and Russian territory, followed the course adopted on several previous occasions, as when the late Dr. Stoliczka was sent as naturalist with the Yarkand mission in 1873-74, and when Dr. J. Anderson was added to the two expeditions to Yunnan in 1868 and 1875. The Indian Government have added to the value of the observations made by publishing the results, which comprise a few general remarks on the fauna, flora and geology, and a descriptive list of the specimens obtained.

The Commission left the valley of Kashmir on June 21, and returned to it on October 12, so that the journey occupied less than four months, whilst the time actually spent on the Pamir itself extended only from July 20 to September 16. This, however, is probably the best time in the whole year for zoological and botanical collecting at so high an elevation, and is certainly much better than April and May, when Stoliczka's collections were made in the same region.

The results, as Mr. Alcock points out, appear small, but this must be attributed to the poverty of the fauna and flora, every effort having been "made to get together as complete and representative a collection as possible." Six mammals, 37 birds, 4 fishes, 10 butterflies, and a few miscellaneous invertebrates were obtained, besides 105 phanerogamous and 10 cryptogamous plants. Four more mammals were seen, but no reptile nor batrachian was met with, despite careful search, and although specimens of both were obtained on the journey between Kashmir and the Pamir. All the fishes, except one loach, belong to the curious group of carps (the *Schizothoracina* of McClelland), with enlarged imbricate scales at the base of the anal fin. This group appears to be peculiar to Central Asia.

In the list of the animals obtained on the road between Kashmir and the Pamir, a very few forms with Indian affinities occur, for instance, a *Trochalopteron* among the birds, but still the great majority are Palearctic species; even three earthworms are identified by Mr. Beddard as European. The Pamir fauna and flora show no trace of Indian affinities, but pertain strictly to the Central Asiatic phase of the Palearctic (or Holarctic) region. At the same time, now that we know the fauna and flora of the Pamir plateau well, their most striking feature is the distinction shown from the animals and plants of Tibet. There is no great difference between the physical features of the two areas; they are both from 12,000 to about 18,000 feet in height (the Tibetan

averaging rather the higher, but by not more than 2000 feet), and the two are completely united by the tableland of Western Tibet. Nevertheless very few of the animals or plants are identical, and the few that are appear to be forms of very wide range. Nor is this all, for so far as the mammalia, the best known and most important group, are concerned, the fauna of the bleak, barren plateau of Tibet appears to be considerably more numerous than that of the rather less bleak Pamir, despite the well-known rich pasture lands of the latter. Thus, taking the Ungulates alone, only two species, *Ovis poli*, Marco Polo's sheep, and *Capra sibirica*, the Asiatic ibex, are known from the Pamir; whilst in Tibet, the yak, Tibetan antelope (*Pantholops*), Tibetan gazelle, two if not three wild sheep (*Ovis hodgsoni*, *O. nahura*, and probably *O. vignei*), an ibex, and the kiang (*Egus hemionus*) occur, without taking into account *Budorcas*, two species of *Cervus* and the musk deer, which are found in parts of the plateau. The same difference is found in other mammalia; thus the golden marmot, *Arctomys aureus*, of the Pamir replaces *A. himalayanus* and *A. robustus* of Tibet, and *Lepus tibetanus* to the westward represents the *L. pallipes* and *L. hypsibius* of the great Eastern plateau.

As already stated the two plateaus, the Pamir and Tibet, are continuous, and are not separated from each other by any distinct elevation or depression. The cause of the marked difference in the fauna and flora needs explanation, and may not improbably be connected with the geological history of the two areas. It has been already shown that the specialisation of the Tibetan mammalian fauna probably indicates isolation during the latter portion of the Tertiary era, an isolation which can only be attributed to elevation. Whether it should be inferred that the elevation of the Pamir, which is believed to have been connected with the origin of the Himalayas, is of later geological date than that of Tibet, is a question that must be left to future geological explorers.

The geological observations in the present work are limited to petrological notes, the rocks found having been noted, and specimens brought away, which were examined by Mr. Holland. This proceeding, like the proverbial carrying away of a brick as a sample of a house, though apparently approved by high geological authorities, is extremely unsatisfactory, and it is to Mr. Holland's credit that he has been able to add one interesting fact, at all events, to previous observations. This is that certain rhyolites which are found on the Pamir are precisely the rocks that might be expected as the volcanic representatives of Stoliczka's "Central Gneiss," which Mr. Holland agrees with General McMahon in regarding as intrusive. This "Central Gneiss" forms the axis of the Himalayas and, as Stoliczka showed, it occupies an extensive area on the Pamir. It is curious that no allusion to Dr. Stoliczka's observations on the Pamir is to be found in Mr. Holland's notes.

In conclusion, it may be fairly stated that we are indebted to the Government of India and to Mr. Alcock for a very useful addition to the facts hitherto known as to the distribution of Asiatic animals and plants. Mr. Alcock acknowledges the assistance given by Mr. Finn in determining the birds collected, and by M. de Nicéville for the Lepidoptera. Mr. Duthie, who supplies the list

of plants, has adopted the admirable plan of giving under each species a brief note of its range, a most valuable addition in a paper of which the importance is chiefly distributional. The plates consist of photo-etchings well executed in the Survey of India Office, and represent fishes, reptiles and crustaceans, the rock structure of a biotite granite, and a view of an *Ovis poli* skin on a wall of rough stones amongst small orchards. The last is so good a plate, that it is impossible to help regretting that a more congenial background has not been selected.

W. T. B.

SOCIOLOGICAL SCIENCE.

Outlines of Sociology. By Lester F. Ward. Pp. xii + 301. (New York: The Macmillan Company, 1898.)

MR. WARD'S little volume, with its clear thought and trenchant writing on more than one topic of current interest, will be welcomed by all students of sociology. It is a reprint of twelve chapters formerly contributed by the author to the *American Journal of Sociology* during the years 1895 to 1897. In the first six lectures, which bear the general title "Social Philosophy," Mr. Ward discusses the old question of the proper position of sociology in a systematic classification of the sciences. The general philosophical position adopted is that of Comte, but the author very properly restores anthropology and psychology to their lawful position in the scheme of the sciences between biology and sociology, and insists with great force upon the very special dependence of sociological on psychological science. The most interesting feature of this part of the book is Mr. Ward's able criticism of Mr. Herbert Spencer's favourite comparison of society to a huge biological organism. Following the lead of Prof. Huxley, he shows, by irresistible arguments, that it is not the whole biological organism, but only the nervous system which really corresponds to a society, and further, that society in its present state is at best a "very low form of organism."

"The most extreme socialist would shrink from the contemplation of any such absolutism as that exercised by the central ganglion of even the lowest of the recognised Metazoa. In order to find a stage comparable to that occupied by society with respect to the central control of the functions of life, it is necessary to go down among the Protozoa and study those peculiar groups of creatures that live in colonies so adapted, that, while the individuals are free to act as they please within certain limits, they are still imperfectly bound together by protoplasmic threads to such an extent that they are in a measure subordinate to the mass thus combined, and really act as a unit or body."

When conscious co-operation of society, as a whole, for its own welfare supersedes sporadic individual effort, and not before will there be a real parallelism between social institutions and the nervous structure of the higher animals.

In the second part of the book, which is entitled "Social Science," Mr. Ward describes the gradual evolution of such a higher form of social structure. Social institutions at first grow up unconsciously under the pressure of the mere "struggle for existence." As intelligence progresses this stage of mere "genesis" passes into the higher stage, called by Mr. Ward "telesis";

unconscious growth gives place to the deliberate manufacture of institutions by conscious purposive action. Hitherto such conscious creation of social institutions has been the work of a few exceptional individuals, but in a higher stage of evolution we may expect it to take the form of "collective telesis," i.e. the deliberate co-operation of the community as an organised whole in the work of social amelioration.

Perhaps the most valuable part of Mr. Ward's book is that in which he discusses the differences between mere unconscious growth and deliberate constructive activity. It has been too much the fashion of sociologists in recent years to argue directly from biological analogies, forgetting that society is at least as much a machine as an organism, and that the presence in all but the lowest stages of social evolution of deliberate human purpose profoundly modifies the whole character of the evolutionary process. As Mr. Ward pithily phrases it, "the environment transforms the animal, but man transforms his environment," a remark which has an obvious bearing upon the application of evolutionary principles to the problems of ethics. Altogether the student who is not content with being told that society "evolves," but wishes to know how specifically social differs from merely biological evolution, will find Mr. Ward's last six chapters singularly luminous and suggestive. The get-up and typography of the book are generally worthy of commendation, but there are some ugly misprints of classical names. A. E. TAYLOR.

OUR BOOK SHELF.

A Text-book of Botany. By Dr. E. Strasburger, Dr. Fritz Noll, Dr. H. Schenck, Dr. A. F. W. Schimper; translated from the German by H. C. Porter, Ph.D. With 594 illustrations, in part coloured. (London: Macmillan and Co., Ltd., 1898.)

THE "Text-book of Botany" issued from the famous institute at Bonn has met with such favour on the part of teachers and students, that it is a matter of surprise that the translation of it into English should have been so long deferred. However it is certain to be extensively used, as the subject is handled from a comprehensive standpoint, and the authors have succeeded in hitting the happy mean between a too elementary and a too advanced treatment.

It is the more to be regretted that, as it was passing through the press, the emendations and corrections which have some time ago appeared in the third German edition were not incorporated in the present volume, which seems based on the first edition in the original language. It is, for example, surprising, and to a student confusing, to find elaborate figures and descriptions of centrospheres in dicotyledonous cells on p. 61, when it is known that the author of this part of the book (Strasburger) has long ago abandoned his belief in their existence, and in the current German text expressly denies their presence in these plants. It may also be doubted whether the book gains at all in value by the somewhat poor coloured illustrations of certain examples of flowering plants, although in this the publishers are but following the original. If, however, they could see their way to reduce the rather high price of the book at the expense of these really useless luxuries, both its own circulation and the temper of the purchaser would improve. For it is not a little remarkable to find a work which in Germany can be bought for 7 marks, costing in its English dress 18s. The book is intrinsically so good that it is to be hoped

that a more moderate price will place it within the means of many students who at present will certainly be debarred from possessing it, save through the intermediation of the second-hand bookseller.

Automobiles sur Rails. By G. Dumont. Pp. 184. (Paris: Gauthier-Villars et Fils. Masson et C^{ie}.)

Régularisation du Mouvement dans les Machines. By L. Lecornu. Pp. 217. (Paris: Gauthier-Villars et Fils. Masson et C^{ie}.)

THESE two volumes belong to the *Encyclopédie scientifique des Aide-Mémoire*, and, like most of the volumes in this series, they contain concise statements of the subjects with which they deal. M. Dumont examines the various systems of horseless traction in use. He begins with steam motors, and then in successive short chapters describes compressed air motors, gas and oil motors, motors driven by carbon dioxide and by ammonia, cable traction, and electric motors. The descriptions are not detailed enough to be entirely satisfactory, nevertheless the volume contains a useful survey of the condition and methods of automobile traction.

M. Lecornu gives in his volume a detailed discussion of the motions of governors of indirect and direct action. His treatment of the various problems involved, and his theorems on the conditions of equilibrium of different governors will interest students of the mathematics and mechanics of machinery.

A Pocket Dictionary of Hygiene. By C. T. Kingzett, F.I.C., and D. Homfray, B.Sc. Pp. 104. (London: Baillière, Tindall, and Cox, 1898.)

THIS pocket-book is intended to be of assistance to medical and sanitary officers in their work, by providing them with concise information upon subjects comprehended in the theory and practice of hygiene. The amount of information given is somewhat unequal, and we should hardly have thought it necessary to include such definitions as: "Adipose, fatty. Anhydrous, without water. Cardiac, pertaining to the heart. Caustic, any substance which destroys animal tissue. Combustion, the process of burning. Emanate, to issue or flow from. Morbid, diseased or unwholesome," &c. Hertz's name is spelt Herz, and Lenard is printed Lennard, in the description of Röntgen rays.

The Secret of the Poles. By Henry Campion. Pp. 48. (Birmingham: White and Pike, Ltd., 1898.)

AMONG the views advanced by the author in this booklet are the following:—The earth is hollow—there is a hollow region large enough to hide the moon and to spare—the earth's axis is hollow—it has two openings, one at each pole—meteoric swarms and ether are attracted through the axis at the south polar entrance, there producing the aurora australis, and after acting as fuel for the fire in the interior is shot out as a waste product at the north polar exit, where it produces the aurora borealis. The character of the book is sufficiently indicated by these extracts, which need no comment.

Wireless Telegraphy, popularly explained. By Richard Kerr, F.G.S. Pp. xv + 111. (London: Seeley and Co., Ltd., 1898.)

MR. W. H. PREECE expresses his general approval of this little volume in a short preface; but at the same time he mentions that he does not accept any responsibility for the controversial points raised. The author explains the principles and practice of telegraphy without intervening wires in words which will be found intelligible by readers unfamiliar with electrical terms. His descriptions possess the merit of being popular in style, and the illustrations assist in brightening the text.

LETTERS TO THE EDITOR

The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, or to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.]

Chance or Vitalism.

PROF. JAPP'S exceedingly interesting address on "Stereochemistry and Vitalism" contains a direct challenge to those who, like myself, hold that we can at present only remain agnostic with regard to the problem of "the possibility or impossibility of living matter originating from dead matter by a purely mechanical process." Unfortunately the facts upon which he bases his view, that chemical compounds of one-sided asymmetry cannot arise save under the influence of life, touch a field so much more familiar to him than to me, that it may seem presumption on my part to make one or two suggestions drawn more directly from my own sphere of work. Still, to the philosophical side of his reasoning I think Clifford, whom he challenges, might, perhaps, have been ready with some reply. I would first state what, I think, stripped of technical language, and represented by the simplest case, is Prof. Japp's standpoint:

(1) Optically active liquids are due to asymmetrical molecules.
(2) These asymmetrical molecules arise from the replacement in a symmetrical molecule of either a right-hand or left-hand atom out of two equal atoms which are images of each other.
(3) No optically active substance can be formed unless there be a selection of purely right-handed or purely left-handed atoms, or, at any rate, unless there be a sensible majority of one or of the other.

(4) Some asymmetric solvents have a power of selective action on optically inactive mixtures of right-handed and left-handed atoms, or, to use the technical term, of two enantiomorphs.

(5) No mechanical process (chemical or physical) could select a right-handed as distinguished from a left-handed atom in a symmetrical molecule, and so produce an asymmetric compound. Any mechanical force which acts on a symmetrical molecule is as likely to affect one atom in a molecule as its image. If I. be the selecting of a right-hand atom and II. of a left-hand atom, then, as Prof. Japp puts it:

"The chances in favour of these two events being equal, the ratio,

Number of occurrences of event I.

Number of occurrences of event II.

will, if we are dealing with an infinitely great number of molecules, approximate to unity. We therefore obtain a mixture, optically inactive by inter-molecular compensation."

Now, putting on one side any objections to Prof. Japp's reasoning arising from the fact that it is based on a purely geometrical hypothesis as to the constitution of molecules, which is also merely descriptive, for we can have no ultimate evidence of its actuality¹—I would draw attention to the fact that (5) is an appeal to the doctrine of chance, and that Prof. Japp tells us that the *mechanical* production of a left-handed or right-handed enantiomorph is an event like the tossing of a coin, of which the chances are precisely equal as to heads or tails. Now if Prof. Japp will toss twenty coins, ten heads and ten tails will undoubtedly be the most frequent result; but there will be a variation about this mean result, and if he goes on tossing long enough he will ultimately come to an instance of twenty heads or twenty tails alone. Cases in which there is a preponderance of heads or tails of a very sensible kind will not be very infrequent. I take it that it is only a *majority* of left- or right-handed enantiomorphs which is required to produce an optically active substance. What majority might be easily ascertained by delicate experiments on the rotatory power of mixtures of dextro- and laevo-acids which are mirror-images of each other. Now, according to Prof. Japp, chance is the factor at work in the production of optically inactive mixtures of right- and left-handed enantiomorphs. Hence, it may be in the course of indefinite ages, purely mechanical action must certainly have produced chemical compounds of one-sided asymmetry with various degrees of rotatory power, due to the greater or less

¹ Such a *geometrical* hypothesis cannot give the *dynamical* explanation of rotatory polarisation required by the physicist, and therefore the "eminent physicist" quoted by Prof. Japp was, I venture to think, right in saying that an explanation of rotatory polarisation is still wanting.

frequency of the two enantiomorphs. We might even predict that if a chemist were to spend his life in the preparation of innumerable and smallest physically sensible amounts of a normally racemoid substance, he would with fine enough apparatus ultimately be able to detect some amount of rotatory polarisation. In nature, where during countless ages (and in the past probably much more actively than at present) inorganic actions and reactions have taken place without man's aid, the production of chemical compounds of one-sided asymmetry must, on Prof. Japp's view of the relation of mechanical action to chance, undoubtedly have taken place. Further, according to Prof. Japp's fourth principle, we may look upon such asymmetrical compounds when they have once arisen as "breeders," or endowed with a power of selecting their own kind of asymmetry from other racemoid substances. Let us put this further statement on one side, however, and content ourselves solely with dissent from Prof. Japp's view expressed in the words:

"But the chance synthesis of the simplest optically active compound from inorganic materials is absolutely inconceivable. So also is the separation of two crystallised enantiomorphs under purely symmetric conditions."

On the contrary, if the theory expounded by Prof. Japp be correct, the inorganic origin of optically active compounds is not only conceivable, but it has a degree of probability which, however small, might be calculated when we know what is the minimum number of molecules in a physically just sensible solution, and what is the majority of enantiomorphs of one kind which will give a just measurable amount of rotatory polarisation.

It will indeed be a great gain if Prof. Japp's address calls more attention to this exciting subject, and leads to further experiment and research.

KARL PEARSON.

The Moon's Course.

THE annual course which the moon takes in company with the earth round the sun was to me a long time a great puzzle, as it is to many others, until one day I demonstrated it to myself by the simplest method. Those who have some smattering of the heavenly bodies generally fail in their attempt to draw the moon's orbit; they find no explanation in popular works, and even in books written by well-known authors the subject of the moon's motion is altogether ignored. All that is found is a circle showing the moon's phases, and it is this circle which is fatal to the conception of the true orbit of the moon: even very young readers see the impossibility of a dozen or thirteen circles surrounding the sun.

For myself, failing to find assistance in books, I readily solved the problem by a practical method. I took a piece of wire and placed a cork on each end. I then drew a line on the table and advanced one, which I called the earth, along the line; the other cork necessarily followed it, and at the same time circulated round the earth in accordance with another force or motion given to it. Beginning with the moon behind the earth and making it pass to the right, both advancing forward, the moon made a curve until it reached the front of the earth, and then the latter still advancing, took the moon with it; this came on its left side, and then making another curve on that side, again reached the front; the two curves completing the orbit round the earth.

My object in writing at the present moment is, that when so many persons are spending their holidays at the sea-side and there is much talk of the tides together with the moon, I take the opportunity of demonstrating by a similar method the moon's course on the sands. I make one person walk in a straight line, marking this with a stick, which he drags behind him, and I call him the earth. I then place another person, the moon, at a given distance from the other, telling him to advance also, and at the same time circulate round his companion. Beginning behind, and taking the right hand, he goes forward, making a curve until he gets to the front, then passes to the left side and forms another curve just as in the other experiment. The four quarters of the moon are in this manner seen, and if the second person continually faces the earth the moon's monthly revolution is also completed.

As these demonstrations are made on a plane, the experiment may also be shown in another way. I wind some wire thirteen times (the number of the lunar months) round a cylinder, and then take off the coil thus made. I pull it out into a helix and

join the ends of the wire. The course of the moon is shown in the spiral, although no correct proportions are attempted. This also gives some idea why the moon is sometimes seen on the horizon and sometimes at the zenith; the reason of which is a great difficulty to young people, as they find nothing about it in their books.

I make no pretence to be an astronomer or mathematician, and, indeed, it may be even a presumption to send this communication to a scientific journal; but if those who can teach fail to do so in popular books, there is no other method for the uninitiated to do the best for themselves.

SAMUEL WILKS.
Grosvenor Street.

The Aurora of September 9.

THIS evening, at about three minutes past eight, on looking out towards the south-west, I was struck by the appearance of a shaft of white light stretching from that direction upwards towards a point 10° or 20° south of the zenith, and immediately on reaching open ground, whence a wide view was obtained, became aware that a fine display of aurora borealis was in progress. The shaft of light successively appeared and disappeared at intervals of a few seconds, and each time further eastwards, but each section was separated from the last by a space of unilluminated sky, as if there were regular spaces in the course of the beam incapable of being set aglow. Very soon after passing overhead the illumination became faint and disappeared eastwards, but now a strong glow appeared again near the horizon under a cloud south-west by west, and this rapidly grew upwards, repeating the phenomenon described in the first case. This occurred at least eleven or twelve times with little variation, except that the shaft became broader and more diffuse. The glow near the horizon south-westwards was always followed by the passage of the light across the sky within 1 to $1\frac{1}{2}$ mins., and each travelling beam was separated from the last by an interval of $1\frac{1}{2}$ to 2 mins. Meanwhile the sky north-westwards showed a pale white steady auroral light, which was at first attributed to the remains of twilight, but which continued to increase till about 8.30, when it was sufficiently strong to cast shadows and to show large print distinctly. This illumination extended slowly upwards from the north until it covered a great part of the sky up to the zenith, and seemed to be due to the luminosity of the great sheet of cirrus and cirrostratus which had covered the sky since 3.30 p.m., for it exactly imitated the distribution forms of the cloud which I had noted during the afternoon. This cloud, unilluminated, did not appreciably obscure the stars. The lacunae of dark interspaces, while the surrounding sky shone with pale light, resembled bands or puffs of dark smoke, but remained fixed in the same situation, and altogether the phenomenon was strangely persistent, the only evident changes being a gradual shifting from north-west to north-east, and at about 9 p.m. a rosy streak north-eastwards. A little later fresh rays were shooting up from north-west by west, somewhat resembling those which an hour previously had emerged from south-west by west, but more steady and unbroken. At 9.55 a faint light was still to be seen in a north-westerly direction. Two hours before this grand exhibition we had been able to get a good view of the large spot on the sun, the intervening bank of cirriform cloud making his appearance through a telescope more like that of the moon. I have not seen any authoritative statements as to interruption of telegraphic messages or other magnetic disturbances coincident with or preceding the visible aurora, but no doubt some effect must have been observed.

Dunrozel, Haslemere, September 9. ROLLO RUSSELL.

A White Sea.

DURING the recent voyage of the P. and O. Co. s.s. *India* to London I had an opportunity (owing to the kindness of Captain Worcester) of witnessing what seems to be a rare phenomenon. The commander had seen it two or three times in the course of his experience. No one else on board knew anything about it, and I should be glad of a reference to any detailed description.

At 1.30 a.m. on the morning of August 22, in the Indian Ocean, the officer in charge saw ahead what seemed to be a low mist, and into which the vessel steamed. I was called about fifteen minutes later.

The whole sea was milk-white, much more luminous than the clear, starry sky, and there was a very definite horizon.

There was no moon, the wind was south-west and light—the end of the monsoon; and although the sea was, as a matter of fact, breaking here and there, it appeared a calm white sheet, only disturbed by the displacement waves near the ship and a very occasional breaker elsewhere; showing through it were occasional flashes of the ordinary brilliant phosphorescence. It will therefore be seen that the luminosity of the “white sea” was rather less than that of a breaking wave with the same illumination. A bucket of water drawn showed nothing unusual. Samples with and without alcohol were preserved.

A fireball was thrown overboard, and burnt on the surface of the water; this was done in order to see if any fog or mist was present. There was no indication of anything of the kind.

On the port side of the ship is an aperture through which the surplus water from the bath tanks is constantly ejected, slightly warmed. This water, as it fell on to the sea, appeared much blacker than the sea, and floated for a few seconds as a black mass; unfortunately, the same shoot is used for the ashes at times. But the ejected water is quite white by daylight.

The appearance of the sea lasted about an hour, then faded, then brightened again, and was quite bright at daylight, 4.15 a.m.; so that it was seen throughout a distance of nearly fifty miles. A slight recurrence was observed the following night, when the monsoon was blowing more strongly.

At 3 o'clock on the 22nd, in the midst of the “white sea,” the latitude was $10^{\circ} 35' N.$ and the longitude $63^{\circ} 25' E.$; the temperature of the air was $77^{\circ} F.$, that of the water $77^{\circ} F.$ Specific gravity of the water by ship's instrument No. 1314 = 25.

I shall be glad to hand over the specimens of water to any one interested.

JAMES W. BARRETT.

22 Cavendish Square, September 13.

Deep-Sea Dredging, and the Phosphorescence of Living Creatures, at Great Sea Depths.

May I call attention to this most interesting subject, upon which so little is known, and with reference to the exploration of the bottom of tropical seas; *nothing is known*, though there is here a mine of natural history wealth probably of unexampled magnitude. In that interesting work on “The Depths of the Sea,” by Sir Wyville Thomson, published more than twenty years ago, we get a glimpse of a hitherto unworked zoological province, which creates a desire to know more from the richness and beauty revealed, where it would be least expected.

He writes—“We had a gorgeous display of luminosity, coming down the Sound of Skye, while dredging in 100 fathoms.

“The Pavonariae came up, resplendent with a pale lilac phosphorescence, like the flame of cyanogen gas—not scintillating, but constant and sufficiently bright to make every portion of a stem distinctly visible, and the stems were a metre long, fringed with hundreds of polyps; and from the number of specimens brought up, we must have passed through a luminous forest of them.

“Among Echinoderms, *Ophiacantha spinulosa* was one of the prevailing forms, and we were greatly struck with the brilliancy of its phosphorescence. Very young *Ophiacantha* shone very brightly also.

“At 344 fathoms, some of our hauls were taken late in the evening, and the tangles were sprinkled over with stars of the most brilliant uranium green. The light was not constant, nor continuous all over the stars, but sometimes it struck out a line of fire all round the disc; flashing or glowing up to the centre; then that would fade, and the whole five rays of *Ophiacantha spinulosa* would light up at the ends, and spread the fire inwards.

“At 557 to 584 fathoms, many of the animals dredged were most brilliantly phosphorescent. In some places, nearly everything brought up seemed to emit light, and the very mud itself was perfectly full of luminous specks. The Pinnatulæ, Virgulariæ, and Gorgoniæ, shone with a lambent white light, so bright that it showed quite distinctly the hour on a watch.

“The light from *Ophiacantha spinulosa* was a brilliant green, coruscating from the centre of the disc, now along one arm, now along another; and vividly illuminating the whole outline of the star-fish.”

From a depth of 567 fathoms, a beautiful scarlet Urchin, *Echinus microstoma*, was obtained. In the year 1846 Keferstein mentions having seen in Stockholm a Crustacean taken from the

depth of 1400 fathoms, of a bright colour. In 1869 and 1870 dredging was carried down to 2435 fathoms by H.M.'s ship *Porcupine*, and the fact that there is an abundant and characteristic invertebrate fauna at that great depth was placed beyond question; but the bottom of the deep sea that has been fairly dredged, may still be reckoned by the square yard; while every haul of the dredge, hitherto used, has brought to light new and unfamiliar forms.

In the number of NATURE for June 30 of this year, there is a most interesting article on deep-sea fishing by means of a trap, an illustration of which is given. These traps are said to have been used at a depth of three thousand fathoms, with complete success. On one occasion a trap that had been lying on the bottom of the Mediterranean, at 700 fathoms depth, for twenty-four hours, brought up 1198 fish, called *Simenchelys parasiticus*. On another occasion, a new crab, one of the largest ever known, *Geryon affinis*, was brought up, and there were sixty-four specimens of it. All this shows how much remains to be done in this province of natural history.

E. L. J. RIDSDALE.

Rottingdean.

The Injection of Cocaine as a Remedy for Stings.

As no one has answered the question asked by Sir J. F. D. Donnelly in your issue of September 8, will you allow me to say that the hypodermic injection of cocaine, or indeed its use in any form, is never quite free from risk. As with most other drugs there is an element of idiosyncrasy, which sometimes produces unexpected and unpleasant results. I believe these occur more frequently, when the drug is injected, than when it is simply applied to the mucous membrane; and when they do occur, they are very alarming. I know that some dentists have given up the practice of injecting cocaine into the gums before extraction on this account, and having been present once when it was done, I should not consent to it again. What we have to remember with regard to the use of all powerful drugs is this, that a few individuals under all conditions, and nearly all under certain conditions, are specially susceptible to their action; and that we cannot determine *a priori* either the individuals so predisposed, or the conditions which render susceptible those not specially predisposed. These drugs must therefore be used with caution, and not be regarded as wholly innocuous.

I should say the hypodermic injection of cocaine into the tongue is undesirable, and only to be done if the pain is intense or the swelling such as to threaten life, in which case it would probably be useful. I would add that so far as I know no solutions of cocaine keep well.

M.D., OXON.

THE GEOGRAPHY OF THE UNITED STATES.¹

THIS volume completes the description of the North American continent with the exception of Mexico; but, although an excellent account of the United States, it leaves North America itself still undescribed. The general reader, of course, will not miss what he has not been trained to expect; and, if he leaves the conception of geography as a science capable of systematic study out of account, the professed geographer will find much valuable material collected with discrimination and stated clearly and modestly. Indeed, except for an implication on the first page that the United States are in many respects more civilised than Canada, the English reader will find nothing to disturb his equanimity even in the account of the revolutionary war or the feats of the *Alabama*. Mr. Gannett writes always as a good American, but is ready to recognise the defects of his country when necessary, and careful to buttress all agreeable statements with statistics which more than prove them.

Of the arrangement of the matter it is impossible to speak with the same satisfaction. The chapters do not flow in the natural sequence desirable in a literary work,

¹ “Stanford's Compendium of Geography and Travel” (New Issue). North America. Vol. II. “The United States.” By Henry Gannett, Chief Geographer of the United States Geological Survey. Maps and Illustrations. Pp. xvi + 466. (London: Edward Stanford, 1898.)

nor are the subjects arranged in alphabetical order usual in a book for occasional reference. It is a little of a shock to turn from mineral resources to population, from great cities to the native Indians, and from commerce to Alaska; even although each chapter in itself is excellent reading.

Mr. Gannett was evidently anxious to resist the temptation of enlarging unduly upon his own special subject, and he has in our opinion gone to the opposite extreme, and lost an opportunity of showing how completely the structure of the country determined by its geology dominates the whole geography of the United States. The introductory chapter does contain a good deal of geology in relation to the configuration of the different natural regions, but the connecting links with the other distributions are wanting. It would be better in a book intended primarily for the general English reader to translate the statistics so freely given into the usual British units of weight and money; the "short tons" for coal and "long tons" for iron-ore are puzzling, and make

establish electric power-houses at once for all needs. So, too, he shows that no country in the world possesses so many ruined cities as the United States—not only the abodes of the early mound-builders and cliff-dwellers, but ruins of yesterday; mushroom towns that teemed with busy thousands in a year, and were abandoned in a month on the failure of a mine or of a company, leaving only "a history of disappointed hopes, of hardships and struggles."

The movements of population are well treated, and a map showing the areas where more than 10 per cent. of the population are foreign-born, and those where more than 10 per cent. of the population are of negro race, displays the interesting fact that the former occupies the whole north and west, the latter the whole south-east, leaving a narrow belt between the two areas. The number of original statistical and physical maps is one of the most striking and satisfactory features of the book, and the illustrations also are admirably selected. At a time when the United States are entering on a new era



FIG. 1.—The Giant Cactus of the Arid Region.



FIG. 2.—Buttes in the Plateau Region.

comparison with other countries difficult. Besides, it would greatly assist the clear conception of such statistics if they were expressed in round numbers.

Apart from these possibilities for improvement, the book contains nothing which we cannot heartily praise. The revision of the text is very thorough, and we have not detected a single erratum.

Many of the topics are handled with surprising freshness, and many interesting points are brought out, such as the changed manner of life of the hardy fishers of the New England coast, who have found an easy and profitable calling as caterers for holiday-makers from the great cities. The author discusses the whole question of American cities, showing how the convenience of the rectangular plan has outweighed æsthetic considerations; and explaining the relative backwardness of the old cities of the east, compared with the new growths of the west, by the vast amount of capital locked up in such archaic conveniences as gas-works and horse or cable cars, while the untrammelled new municipalities can

of their national life, the publication of so accurate and impartial an account of that great country by one of its own citizens is peculiarly appropriate, and deserves a cordial welcome.

HUGH ROBERT MILL.

THE BRITISH ASSOCIATION.

THE concluding meeting of the British Association was held on Wednesday, September 14. Sir William Crookes occupied the chair, and the Mayor of Bristol (Sir R. H. Symes) and the High Sheriff were present, as well as the principal officers and members of the Association. The following report of the meeting is from the *Times* :—

The proceedings were opened by the announcement that the general committee had been able to pass grants to the amount of 1485*l.*—an amount which was justified largely by the success of the present meeting.

Sir Norman Lockyer proposed that the thanks of the Association be given to the Mayor, the High Sheriff, the

executive committee, and the local officers. He said that the great success of the meeting was largely due to the efforts of those referred to in the resolution. Never before in his experience of the Association had local effort led to such absolute smoothness in the working of the machine. It was a pity that the work of the Association had been so hard as to prevent many members from seeing all the points of interest in this interesting city of Bristol. The magnificent educational establishments which crowded the city were themselves worthy of close attention, and at some future meeting the British Association might find Bristol at the head of some great southwestern University.

Prof. Schäfer seconded the resolution.

Mr. James Scott (of Toronto), on behalf of the Canadian members, expressed high appreciation of the welcome which had been accorded to them.

The resolution was then carried with much enthusiasm.

The Mayor of Bristol, Mr. Howell Davis (chairman of the executive committee), Mr. Arrowsmith (local treasurer), and Mr. Arthur Lee and Dr. Bertram Rogers (local secretaries), each responded, Mr. Arrowsmith expressing his acknowledgments for the cheque for 120*l.* which had been given by the Council of the Association towards the Colston Hall fund.

Prof. Rücker next moved a comprehensive vote of thanks to all public bodies and private persons who had contributed to the success of the meeting. He said that if the citizens of Bristol had not supported the local officers, the success of the meeting could not have been secured in so large a measure. As President of the International Committee of the Magnetic Conference, he was charged to convey the best thanks of the foreign members to the Association and to the local authorities for the extreme kindness of their reception.

Dr. Gladstone seconded the resolution, which was carried unanimously, the High Sheriff responding.

Sir John Evans moved a cordial vote of thanks to Sir William Crookes, President, for his admirable address and for his conduct in the chair. He prophesied, when introducing Sir William to the chair, that the Association would hear from him a remarkable address, and that prophecy had been amply justified. One of the most valuable portions of that address was that in which public attention was called to the fact that there was in our atmosphere an inexhaustible supply of nitrogen, and that means should be discovered for employing that nitrogen to increase the produce of the earth. Sir William Crookes had fulfilled with courtesy and distinction all the many and various duties which the past week had imposed upon him, and the thanks of the Association were cordially due to him.

Prof. Roberts-Austen seconded the resolution, which was carried with enthusiasm.

Sir W. Crookes, in responding, said that he felt like an electrical switch-board—for really he was only the transmitter and distributor of these thanks to those whose help had been so material. He was especially grateful to the Mayor and Mayoress, whose hospitality had facilitated his work so greatly; and he regarded as one of the highest compliments ever paid to him the invitation to the remarkable smoking-symposium of the previous Friday evening. As for the President's office, the pace was getting too fast for human endurance; and in a short time the British Association would, if the work were to be got through at all, have to elect a young athletic man of five-and-twenty instead of a man over three-score years and ten.

Prof. Rücker announced that the number of tickets issued for the present meeting of the Association was 2446.

This concluded the proceedings. The next meeting will be held at Dover, and will commence on September 13, 1899.

SECTION D.

ZOOLOGY.

OPENING ADDRESS BY PROF. W. F. R. WELDON, M.A., F.R.S., PRESIDENT OF THE SECTION.

IN attempting to choose the subject of the address with which custom obliges your president to trouble you, I felt that I should have the best hope of interesting you if I decided to speak to you on the subject most interesting to myself. I therefore propose to discuss, as well as I can, the principal objections which are urged against the theory of Natural Selection, and to describe the way in which I think these objections may be met.

The theory of Natural Selection is a theory of the importance of differences between individual animals. In the form in which Darwin stated it, the theory asserts that the smallest observable variation may affect an animal's chance of survival, and it further asserts that the magnitude of such variations, and the frequency with which they occur, is governed by the law of chance.

Three principal objections are constantly brought forward against this theory. The first is that the species of animals which we know fall into orderly series, and that purely fortuitous variations cannot be supposed to afford opportunity for the selection of such orderly series; so that many persons feel that if the existing animals are the result of selection among the variable offspring of ancestral creatures, the variations on which the process of Natural Selection had to act must have been produced by something which was not chance.

The second objection is that minute structural variations cannot in fact be supposed to affect the death-rate so much as the theory requires that they should. And it is especially urged that many of the characters, by which species are distinguished, appear to us so small and useless that they cannot be supposed to affect the chance of survival at all.

The third objection is that the process of evolution by Natural Selection is so slow that the time required for its operation is longer than the extreme limit of time given by estimates of the age of the earth.

Now the first of these three objections, the objection to fortuitous variation as the source of material on which Natural Selection can act, is very largely due to a misunderstanding of the meaning of words. The meaning of the word Chance is so thoroughly misunderstood by a number of writers on evolution that I make no apology for asking you to consider what it does mean.

Consider a case of an event which happens by chance. Suppose I toss a penny, and let it fall on the table. You will agree that the face of the penny which looks upwards is determined by chance, and that with a symmetrical penny it is an even chance whether the "head" face or the "tail" face lies uppermost. For the moment, that is all one can say about the result. Now compare this with the statements we can make about other moving bodies. You will find it stated, in any almanac, that there will be a total eclipse of the moon on December 27, and that the eclipse will become total at Greenwich at 10.57 p.m.; and I imagine you will all feel sure, on reading that statement, that when December 27 comes the eclipse will occur; and it will become total at 10.57 p.m. It will not become total at 10.50 p.m., and it will not wait until 11.0 p.m. You will say, therefore, that eclipses of the moon do not occur by chance.

What is the difference between these two events, of which we say that one happens by chance, and the other does not? The difference is simply a difference of degree in our knowledge of the conditions. The laws of motion are as true of moving pence as they are of moving planets; but it happens that we know so much about the sun, and the earth, and the moon, that we know the circumstances which affect their relative positions very accurately indeed; so that we can predict within less than a minute the time at which the shadow of the earth will next fall upon the moon.

But the result of tossing a penny depends upon a very large number of things which we do not know. It depends on the shape and mass of the penny, its velocity and direction when it leaves one's hand, its rate of rotation, the distance of one's hand from the table, and so on. If we knew all these things before tossing the penny, we should be able to predict in each case what the result would be, and we should cease to regard pitch and toss as a game of chance.

As it is, all we know about these complicated conditions is that if we toss a penny for a number of times, the conditions

which give "heads" will occur about as often as the conditions which give "tails."

If you examine any event which occurs by chance, you will find that the fortuitous character of its occurrence always depends upon our ignorance concerning it.

If we know so little about a group of events that we cannot predict the result of a single observation, although we can predict the result of a long series of observations, we say that these events occur by chance. And this statement seems to me to contain the best definition of chance that can be offered.

If we used the word chance in this sense, we see at once that our knowledge of animal variations is precisely knowledge of the kind referred to in our definition of chance. We know with some certainty the average characters of many species of animals; but we do not know exactly the character of the next individual of these species we may happen to look at. So that in the present state of our knowledge it is *a priori* certain that the great majority of animal variations should occur by chance, in the sense in which we have used the phrase; and I will show you in a moment illustrations of the fact that they do so occur.

But before doing so, I would point out the difference between the sense in which we have used the word chance, and the sense in which it is used by many objectors to the theory of Natural Selection. Such epithets as *blind*, *lawless*, and the like, are constantly applied to chance; and a kind of antithesis is established between events which happen by chance, and those which happen in obedience to natural laws. In many German writings, especially, this antithesis between *Zufälligkeit* and *Gesetzmässigkeit* is strongly insisted upon, whenever organic variation is discussed.

This view of chance is not supported by experience; and indeed, if it could be shown that any thing in human experience were absolutely lawless, if it could be shown that in any department of nature similar conditions did not produce similar effects, the whole fabric of human knowledge would crumble into chaos, and all intellectual effort would be a profitless waste of time. There is not the slightest reason to believe that any such absolutely lawless phenomena do exist in nature; so that we need pay no further attention to the writers who assume that chance is a lawless thing.

But if chance is a perfectly orderly and regular phenomenon, then the question, whether animal variations occur by chance or not, can be settled by direct observation. I will now show you one or two examples of events which undoubtedly occur by chance, and then compare these with one or two cases of organic variation.

As events which occur by chance, I have taken the results of tossing twelve dice. My wife has spent some time during the last two months in tossing dice for you, and I will ask you to look at the results.

Her first record gives the number of dice showing more than three points in each of 4096 throws of twelve dice. There are, of course, six numbers on each of the dice; so that if all the dice were perfectly symmetrical and similar, the average number of dice with more than three points should be six in each throw of twelve. But dice are not symmetrical and similar. The points on the dice used were marked by little holes, scooped out of their faces; and the face with six such holes scooped out of it was opposite to the face with only one such hole: so that the face with one point was heavier than the face with six points; and therefore six was rather more likely to be uppermost than one. In the same way, two was opposite five; so that the five face was a little more likely to fall uppermost than the face with two points. Therefore, it is a little more likely that you will throw four, five, or six, in throwing dice, than it is that you will throw one, two, or three.

Accordingly, the average number of dice, in these 4096 throws, which had more than three points, was not six, but 6.135.

To show you that this excess of high points was due to some permanent property of the dice, she threw these twelve dice another 4096 times; and the average number of dice with more than three points was 6.139. A third series of trials gave an average of 6.104, and a fourth gave an average of 6.116.

You see that the difference between the highest and the lowest of these determinations is only about one-half per cent., so that the mean result of such a series of fortuitous events can be determined with great accuracy.

And just as the mean of the whole series can be determined, so we can know with considerable accuracy how often any

possible deviation from the average result will occur. The degree of accuracy with which we can know this may be judged from Table I.

TABLE I.—Frequency with which Dice showing more than three Points were thrown in each of Four Series of Trials, the number of throws in each Series being $2^{12} = 4096$.

Number of dice with more than 3 points.	Most probable frequency for symmetrical dice.	Observed frequencies.			
		I.	II.	III.	IV.
12	1	0	1	0	1
11	12	11	13	8	14
10	66	71	86	61	66
9	220	257	246	241	241
8	495	536	540	513	586
7	792	847	836	856	861
6	924	948	913	948	866
5	792	731	750	802	728
4	495	430	446	420	474
3	220	198	198	182	204
2	66	60	55	51	67
1	12	7	12	13	6
0	1	0	0	1	0

You see that the results of the experiments agree fairly well with one another, and differ from the results most probable with symmetrical dice, in the way which the structure of the actual dice would lead one to expect. Throws which give seven, eight, or nine dice with more than three points occur too often, throws in which only two, three, or four dice have more than three points do not occur often enough. You see then that each of these results is orderly and regular, and that the four results agree very fairly among themselves, not only in the mean value of each of them, but in the magnitude and frequency of departures from the mean. That they differ from the results which would probably be obtained with symmetrical and similar dice is only to be expected, because the dice used are neither symmetrical nor similar.

You notice that this table is very nearly symmetrical; the most frequent result is that which lies in the middle of the series of possible results; and the other frequencies would, with perfect dice, be distributed symmetrically on each side of it; so that with perfect dice one would be as likely to throw five dice out of twelve with more than three points as one would be to throw seven, and so on.

This symmetry in the distribution of the results is only found when the chance of the event occurring in one trial is even. The next table shows the result of 4096 throws of twelve dice.

TABLE II.—Frequency of Sixes in 4096 throws of Twelve Dice.

Number of sixes.	Most probable number with symmetrical dice.	Number observed.
8	0.58	1
7	4.66	7
6	27.18	24
5	116.43	115
4	363.84	380
3	808.53	796
2	1211.44	1181
1	1102.56	1145
0	459.52	447

in which sixes only were counted. The chance against throwing six with any one of the dice is of course five to one; so that in throwing twelve dice you are more likely to throw two sixes than to throw any other number. But you see that the chance of throwing only one six is very much greater than the chance of throwing three; the chance of throwing none is greater than the chance of throwing four, and while there is a chance of throwing five, six, or more, of course it is impossible to throw less than none at all; so that the diagram is all askew. You see that this time, as before, the frequency with which any

number of sixes did actually occur was as near to the result most probably with perfect dice as the asymmetry of the actual dice allows one to expect.¹

These results will be enough to show you how absurd is the attitude which so many writers have taken up towards chance when discussing animal variation. The assertion that organic variation occurs by chance is simply the assertion that it obeys a law of the same kind as that which expresses the orderly series of results we have just looked at.²

That is a matter which can be settled by direct observation. But in order to express the law of chance in such a way that we can apply it to animal variation, we must make use of a trick which mathematicians have invented for that purpose.

It is a well-known proposition in probability that the frequency with which one throws a given number of sixes in a series of trials with twelve dice is proportional to the proper term in the expansion of $(\frac{1}{2} + \frac{1}{2})^{12}$. The values in this table were calculated by expanding this expression. But if I had wanted to show you the most probable result of experiments with 100 dice, I should not willingly have expanded $(\frac{1}{2} + \frac{1}{2})^{100}$. The labour would be too enormous. Then again, suppose we are given a number of results, and are not told how many dice were used, how are we to find out the power to which we must raise $(\frac{1}{2} + \frac{1}{2})$, since this depends on the number of dice?

Before applying the law of chance to variations in which we cannot directly measure the number of contributory causes (the analogue of the number of dice), we must find some way out of this difficulty.

The way is shown by the diagram (Fig. 1).

The rectangles in this diagram are proportional to the various terms of $(\frac{1}{2} + \frac{1}{2})^{12}$; and they represent the most probable result of counting the number of dice with more than three points in a series of trials with twelve dice. The heights of these rectangles were determined by expanding $(\frac{1}{2} + \frac{1}{2})^{12}$; but you notice the dotted curve which is drawn through the tops of them. The general slope of this curve is, you see, the same as the general slope of the series of rectangles; and the area of any strip of the curve which is bounded by the sides of a rectangle is very nearly indeed the same as that of the rectangle itself.

The constants upon which the shape of this curve depends are easily and quickly obtained from any series of observations; so that you can easily and quickly see whether a set of observed phenomena obeys the symmetrical law of chance or not.

A good many characters of animals do vary in this symmetrical way; and I show you one, which will always be historically interesting, because it was one of the principal characters used to illustrate Mr. Galton's invaluable applications of the law of chance to biological problems. That is the case of human stature. The diagram (Fig. 2) shows the stature of 25,878 American recruits; and you see that the frequency with which each stature occurs is very close indeed to that indicated by the curve. So that variations in human stature do occur by chance, and they occur in such a way that variation in either direction is equally probable.

In cases where a variation in either direction is equally likely to occur, this symmetrical curve can be used to express the law

of distribution of variations. And the great difficulty in applying the law of chance to the treatment of other cases was, until quite lately, that the way of expressing asymmetrical distributions by a similar curve was unknown; so that there was no obvious way of determining whether these asymmetrical distributions obeyed the law of chance or not.

The form of the curve, related to an asymmetrical distribution of chances, as the curve before you is related to symmetrical distributions, was first investigated by my friend and colleague Prof. Karl Pearson. In 1895 Prof. Pearson published an account of asymmetrical curves of this kind, and he showed the way in which these curves might be applied to practical statistics. He illustrated his remarkable memoir by showing that several cases of organic variation could be easily formulated by the method he described; and in this way he made it possible to apply the theory of chance to an enormous mass of material, which no one had previously been able to reduce to an orderly and intelligible form.

In this same memoir Prof. Pearson dealt with another




FIG. 1.

problem in the theory of chance, which has special importance in relation to biological statistics. It has doubtless occurred to many of you that the analogy between the complexity of the results obtained by tossing dice, and the complexity of events which determine the character of an animal body, is false in an important respect. For the events which determine the result, when we throw a dozen dice on the table, affect each of the dice separately; so that if we know that one of the dice shows six points, we have no more reason to suppose that another will show six points than we had before looking at the first.³ But the events which determine the size or shape of an organ in an animal are probably not independent in this way. Probably when one event has happened, tending to increase the size of an arm or a leg in an embryo, it is more likely than it was before that other events will happen leading to increased size of this arm or leg. So that the chances of variation in the size of a limb would be represented by a law similar to that which expresses the result of throwing dice, but different from it. They would more nearly resemble the result of drawing cards out of a pack. Suppose you draw a card out of a pack. It is an even

³ That is to say, if we know beforehand that the dice are symmetrical.

¹ It is unfortunate that I chose dice as instruments in these experiments. Dice are not only sensibly asymmetrical, but any ordinary dice are sensibly dissimilar; so that the result most probable with any actual dice is not given by a simple binomial expansion. The result theoretically most probable for the actual dice used could not be determined without very careful measurement of the dice themselves; and I was unable to attempt measures of the requisite accuracy. All that the records show, as they stand, is the amount of agreement between four successive observations of a fortuitous event.

² The law is not, however, identical in the two cases; see *infra*.

chance whether you draw a red card or a black one. Suppose you draw a red card, and keep it. The chance that your second card will be red is not so great as the chance that it will be black; because there are only twenty-five red cards and twenty-six black cards left in the pack.

Now Prof. Pearson has shown how to deal with cases of this kind also; and how to determine, from the results of statistical observation, whether one is dealing with such cases or not.

I am no mathematician, and I do not dare even to praise the mathematical process by which this result was achieved. I will only say that it is experimentally justified by the fact that most statistics relating to organic variation are most accurately represented by the curve of frequency which Prof. Pearson deduces for the case where the contributory causes are mutually interdependent.¹

The first case of an asymmetrical distribution in animals which I ask you to look at is the frequency of variations in the size of part of the carapace of shore crabs. The crabs measured were 999 females from the Bay of Naples. In this case the distribu-

series of deviations from the mean length of the antero-lateral margin is as definite a character of the crabs as the mean itself; and in every generation a series of deviations from the mean is regularly produced, according to a law which we can learn if we choose to learn it.

Now suppose it became advantageous to the crabs, from some change in themselves or in their surroundings, that this part of their carapace should be as long as possible. Suppose the crabs in which it was shorter had a smaller chance of living, and of reproducing, than the crabs in which it was longer.

Suppose that crabs in which this dimension is longest were as much more productive than those in which it was shortest, as the most prolific marriages are more fertile than the least prolific marriages among ourselves. Prof. Pearson has pointed out that half the children born in England are the offspring of a quarter of the marriages. If we suppose the productiveness among crabs to vary as much as it does among ourselves, only that in crabs the productiveness is greater, the greater the length of this bit of the carapace, then half of the next




FIG. 2.—Diagram showing the height (in inches) of each of 25,878 American recruits.

tion of variations (see Fig. 3) is very nearly symmetrical, and in an account of these crabs which I wrote before Prof. Pearson's memoir was published, I treated them as symmetrical. The curve actually drawn on the diagram is one constructed by Prof. Pearson himself from the data given by my measurements of the crabs, and it fits the observations very sensibly better than the symmetrical curve. So that this dimension of a crab's carapace does vary by chance, but the chance of a given deviation from the mean length is not quite the same in both directions.

Now, admitting for the moment that these differences in the length of a part of the crab's carapace can affect the crab's chances of survival, you see that natural selection has abundant material on which to work. The production of this regular

generation of crabs will be produced by that quarter of the present generation in which the antero-lateral margin is longest. And as the offspring will inherit a large percentage of the parental character, the mean of the race may be sensibly raised in a single generation.

This view of the possible effect of selection seems to have escaped the notice of those who consider that favourable variations are of necessity rare, and likely to be swamped by intercrossing when they do occur. You see that in this case there are a few individuals considerably different from the mean in either direction, and a very large number which differ from the mean a little in either direction. If such deviation be associated with some advantage to the crabs, so that crabs which possess such abnormality are more fertile than those which do not, it is a certainty that the mean character of the next generation will change, if only a little, in the direction advantageous to the race; and the opportunity for selective modification of this kind to occur in either direction is very nearly the same.

¹ Even the distribution of human stature, which has been so successfully treated by the older, so-called "normal" curve, is more accurately represented by a curve of Prof. Pearson's type; but in this case the difference between the two is so slight as to be inappreciable for all practical purposes; so that Mr. Galton's practice and Prof. Pearson's theory are alike justified.

In the next case, this is not true.

The diagram (Fig. 4) represents the number of female swine, out of a batch of two thousand examined in Chicago, which have a given number of Müllerian glands in the right fore-leg.

amount of possible change is greater in one direction than in the other.

Now let us pass on to another example.

Table III. shows the variation in the number of petals in a race of buttercups studied by Prof. de Vries. You see that the most frequent




FIG. 3.—Diagram showing the magnitude of the antero-lateral margin (in terms of carapace-length) in 999 female shore-crabs from Naples.

The distribution is much more skew than in the case of the crabs, and you see again the very beautiful way in which Prof. Pearson's curve expresses it. You see that the range of variation is much greater on one side of the mean than on the other; and the selective destruction necessary in order to raise the mean number of glands by one would be very different from the amount of destruction necessary in order to lower the mean by one. Further, the mean number of glands in these pigs is $3\frac{1}{2}$; the number which occurs oftenest, the "modal" number as Prof. Pearson calls it,¹ is three. Now it is impossible to lower this number till it is less than 0, so that it can only be diminished by three; but it is conceivable that it should be increased by more than three. So that the amount of selective destruction required in order to change either the mean or the modal character of these pigs in one direction, would be greater than the amount required, in order to produce a change of equal magnitude in the opposite direction, and the

¹ All attempts to confine the word "average" to the most frequently occurring magnitude, and the word "mean" to the arithmetic mean of the series, have failed to secure support. Therefore Prof. Pearson's proposal to call the value which occurs oftenest the "mode" is very useful.

have less than five petals, though a considerable number have more than five; and here again you see the way in which Prof. Pearson's formula fits the observations.

TABLE III.—Prof. Pearson's expression for the variation in the race of Buttercups described by Prof. de Vries.

No. of petals	11	10	9	8	7	6	5	4	3	2	1	0
Observed frequency	11	10	9	8	7	6	5	4	3	2	1	0
Pearson's theory	11.0	10.8	9.6	7.6	23	22.6	13.3	13.6	9.5	5.5	1.3	0.2




FIG. 4.—Diagram showing the number of Müller's glands in each of 2000 female swine.

You see that if this diagram (which is based on very few specimens) really represents the law of variability in these buttercups, no amount of natural or other selection can produce a race with less than five petals out of them. While it is conceivable that selection might quickly raise the normal number of petals, it could not diminish it, unless the variability of the race should first change.¹

These examples, which are typical of others, must suffice to show the way in which the theory of Chance, as developed by Prof. Pearson, can express the facts of organic variation.

I think you will agree that they also show the importance of investigating these facts. For of the four characters we have examined, we have seen that two, namely human stature and the antero-lateral carapace length of *Carcinus menas*, vary so as to afford nearly equal material for selective modification in either direction; one character, the number of Müller's glands in swine, offers distinctly greater facility for selective modification in one direction than in the opposite direction; and in the last character, the number of petals in a race of buttercups appears to offer scope for modification in one direction only, at least by selection in one generation.

Knowledge of this kind is of fundamental importance to the theory of Natural Selection. You have seen that the new method given to us by Prof. Pearson affords a means of expressing such knowledge in a simple and intelligible form; and I, at least, feel very strongly that it is the duty of students of animal evolution to use the new and powerful engine which Prof. Pearson has provided, and to accumulate this kind of knowledge in a large number of cases.

I know that there are people who regard the mode of treatment which I have tried to describe as merely a way of saying, with a pompous parade of arithmetic, something one knew before. This criticism of Prof. Pearson's work was actually made to me the other day by an eminent biologist, whose name I will not repeat. If there be any here who hold such an opinion, I would ask them to read Mr. Francis Galton's Essays on Heredity; where a simple and quite unexpected relation between parents and offspring is shown to be a direct consequence of the fact that they vary by chance. This is the first and the most striking deduction from the mathematical theory of organic variation, but it is not the only one. It is enough, however, to show that the new method is not only a simple means of describing the facts of variation, which facts very few people knew before, but it is a powerful instrument of research, which ought to be quickly and generally adopted by those who care for the problems of animal evolution.

I think I have said enough to convince you how entirely Prof. Pearson's method promises to confirm the assertion that organic variation obeys the law of chance.

The other objections to Darwin's theory are not so easily answered. It is said that small variations cannot be supposed to affect an animal's chance of life or death; but few persons have taken any pains to find out in any given case whether the death-rate is in fact affected by small variations or not. It is said that the process of Natural Selection is so slow that the age of the earth does not give time for it to operate, but I know of few cases in which any attempt has been made to find out by actual observation how fast a species is really changing.

I can only attempt to discuss the importance of small variations, and the rate of organic change, in the one case which I happen to know. The particular case I have myself studied is the variation in the frontal breadth of *Carcinus menas*.²

During the last six years my friend, Mr. Herbert Thompson, and I have studied in some detail the state of this character in the small shore-crabs which swarm on the beach below the laboratory of the Marine Biological Association at Plymouth.

I will show you that in these crabs small changes in the size of the frontal breadth do, under certain circumstances, affect the death-rate, and that the mean frontal breadth among this race of crabs is, in fact, changing at a rate sufficiently rapid for all the requirements of a theory of evolution.

In Table IV. you see three determinations of the mean frontal breadth of these crabs, expressed in terms of the carapace-length taken as 1000. You see that the mean breadth varies very

rapidly with the length of the crab, so that it was necessary to determine it separately in small groups of crabs, such that the length of no two crabs in a group differed by more than a fifth of a millimetre. The first column of the table shows you the mean frontal breadth of twenty-five such groups, between 10 and 15 millimetres long, collected in 1893. These crabs were measured by Mr. Thompson. The second column shows you the mean frontal breadth in twenty-five similar groups of crabs, collected in 1895, and also measured by Mr. Thompson. You see that in every case the mean breadth in a group of crabs collected in 1895 is less than it was in crabs of the same size collected in 1893. The third column contains the result, so far as it is yet obtained, of my own measurement of crabs collected this year. It is very incomplete, because the 1895 crabs were collected in August and September, and I was anxious to compare them with crabs collected this year at the same season, so that there has not yet been time to measure the whole series. The measurements are sufficient, however, to show that the same kind of change has taken place during the last three years as that observed by Mr. Thompson in the interval between 1893 and 1895. Making every allowance for the smallness of the numbers so far measured this year, there is no doubt whatever that the mean frontal breadth of crabs from this piece of shore is considerably less now than it was in 1895 among crabs of the same size.¹

TABLE IV.—The Mean Frontal Breadth ratio of Male *Carcinus menas* from a particular patch of beach in Plymouth, in the years 1893, 1895, and 1898.

Length of carapace.	Mean frontal breadth in terms of carapace length = 1000.				No. of crabs in the 1898 group.
	1893 (Thompson).	1895 (Thompson).	1898 (Weldon).		
10.1	816.17	809.08	—	—	—
10.3	812.06	804.82	—	—	—
10.5	807.37	803.27	—	—	—
10.7	808.96	801.69	—	—	—
10.9	805.07	799.27	—	—	—
11.1	802.50	794.12	784.25	4	—
11.3	798.18	792.38	787.36	11	—
11.5	797.19	788.83	784.00	9	—
11.7	794.28	785.29	782.44	16	—
11.9	791.45	786.53	780.09	11	—
12.1	788.38	780.61	775.25	16	—
12.3	783.98	779.50	773.42	12	—
12.5	783.69	776.50	767.00	11	—
12.7	783.58	773.43	772.43	14	—
12.9	777.38	773.63	764.67	15	—
13.1	776.63	771.61	760.13	16	—
13.3	774.60	766.21	761.29	7	—
13.5	766.91	763.96	759.56	16	—
13.7	767.63	762.00	757.00	16	—
13.9	763.73	759.40	756.10	10	—
14.1	758.94	757.00	742.00	13	—
14.3	756.90	755.77	747.86	7	—
14.5	762.60	754.45	744.44	9	—
14.7	753.00	749.84	739.22	8	—
14.9	751.32	748.03	742.83	6	—

These results all relate to male crabs. The change in female crabs during this time has been less than the change in male crabs, but it is, so far as my measurements at present permit me to speak, going on in the same direction as the change in male crabs.

I think there can be no doubt, therefore, that the frontal breadth of these crabs is diminishing year by year at a rate which is very rapid, compared with the rate at which animal evolution is commonly supposed to progress.

I will ask your patience for a little while longer, that I may tell you why I feel confident that this change is due to a selective

¹ I shall, of course, consider it my duty to justify this statement by more extensive measurement as soon as possible. In the meantime I may say that I have measured other small groups of crabs, male and female, from the same place, at different seasons of the years 1896-98, and the results agree with those recorded in the table.

² Of course we know that selection does change the variability of a race. In 1894 I gave an account of the variation of this dimension in female specimens of various sizes (*Rep. Soc. Proc.*, vol. lvi.). and I put forward an hypothesis of the amount of selective destruction due to variation in this character. That hypothesis neglected several important facts which I now know, and was open to other objections. I desire to replace it by the results of the observations here recorded.

destruction, caused by certain rapidly changing conditions of Plymouth Sound.

If you look at the chart, you will see that Plymouth Sound is largely blocked up, and its communication with the sea is narrowed by a huge artificial breakwater, about a mile long, so that the tidal currents enter it and leave it only by two openings. This huge modern barrier has largely changed the physical conditions of the Sound.

On either side of Plymouth itself a considerable estuary opens into the Sound, and each of these estuaries brings down water from the high granite moorlands, where there are rich deposits of china clay. Those of you who know Dartmoor will remember that in rainy weather a great deal of china clay is washed into the brooks and rivers, so that the water frequently looks white and opaque, like milk. Much of this finely divided china clay is carried down to the sea; and one effect of the breakwater has been to increase the quantity of this fine silt which settles in the Sound itself, instead of being swept out by the scour of the tide and the waves of severe storms.

So that the quantity of fine mud on the shores and on the bottom of the Sound is greater than it used to be, and is constantly increasing.

But this is not all. During the forty or fifty years which have gone by since the breakwater was completed, the towns on the shores have largely increased their population; the great dockyard at Devonport has increased in size and in activity; and the ships which visit the Sound are larger and more numerous than they were. Now the sewage and other refuse from these great and growing towns and dockyards, and from all these ships, is thrown into the Sound; so that while it is more difficult than it used to be for fine silt to be washed out of the Sound, the quantity thrown into it is much greater than it was, and is becoming greater every day.

It is well known that these changes in the physical conditions of the Sound have been accompanied by the disappearance of animals which used to live in it, but which are now found only outside the area affected by the breakwater.

These considerations induced me to try the experiment of keeping crabs in water containing fine mud in suspension, in order to see whether a selective destruction occurred under these circumstances or not. For this purpose, crabs were collected and placed in a large vessel of sea-water, in which a considerable quantity of very fine china clay was suspended. The clay was prevented from settling by a slowly moving automatic agitator; and the crabs were kept in under these conditions for various periods of time. At the end of each experiment the dead were separated from the living, and both were measured.

In every case in which this experiment was performed with china clay as fine as that brought down by the rivers, or nearly so, the crabs which died were on the whole distinctly broader than the crabs which lived through the experiment, so that a crab's chance of survival could be measured by its frontal breadth.

When the experiment was performed with coarser clay than this, the death-rate was smaller, and was not selective.

I will rapidly show you the results of one or two experiments. The diagram (Fig. 5) shows the distribution of frontal breadths, about the average proper to their length, in 248 male crabs treated in one experiment. Of these crabs, 154 died during the experiment, and 94 survived. The distribution of frontal breadths in the survivors is shown by the lower curve in the diagram, and you see that the mean of the survivors is clearly below the mean of the original series, the mean of the dead being above the original mean.

Two other cases, which are only examples of a series in my possession, show precisely the same thing.¹

These experiments seemed to me to show that very finely

¹ It is impossible in this place to give a full account of the experiments referred to, and a multiplication of mere small scale diagrams seems useless, so that only one of those exhibited when the address was delivered is here reproduced.

divided china clay does kill crabs in such a way that those in which the frontal breadth is greatest die first, those in which it is less live longer. The destruction is selective, and tends to lower the mean frontal breadth of the crabs subjected to its action. It seemed to me that the finer the particles used in the experiments, that is to say, the more nearly they approached the fineness of the actual silt on the beach, the more selective their action was.

I therefore went down to the beach, where the crabs live, and looked at the silt there. This beach is made of moderately small pieces of mountain limestone, which are angular and little worn by water. The pieces of limestone are covered at low tide with a thin layer of very fine mud, which is much finer than the china clay I had used in my experiments, and remains suspended in still water for some time. Under these stones the crabs live, and the least disturbance of these stones raises a cloud of very fine mud in the pools of water under them. By washing the stones of the beach in a bucket of sea water, I collected a quantity of this very fine mud, and used it in a fresh series of experiments, precisely as I had before used china clay, and I obtained the same result. The mean frontal breadth of the survivors was always smaller than the mean frontal breadth of the dead.

I think, therefore, that Mr. Thompson's work, and my own, have demonstrated two facts about these crabs; the first is that their mean frontal breadth is diminishing year by year at a measurable rate, which is more rapid in males than in females; the second is that this diminution in the frontal breadth occurs in the presence of a material, namely, fine mud, which is increasing in amount, and which can be shown experimentally to




FIG. 5.—Diagram showing the effect of china clay upon 248 male crabs. The upper curve shows the distribution of frontal breadths in all these crabs; the dotted curve the distribution of frontal breadths in the survivors. The dotted line s shows the mean of the survivors; the line d the mean of the dead.

destroy broad-fronted crabs at a greater rate than crabs with narrower frontal margins.

I see no shadow of reason for refusing to believe that the action of mud upon the beach is the same as that in an experimental aquarium; and if we believe this, I see no escape from the conclusion that we have here a case of Natural Selection acting with great rapidity because of the rapidity with which the conditions of life are changing.

Now, if we suppose that mud on the beach has the same effect upon crabs as mud in an aquarium has, we must suppose that every time this mud is stirred up by the water, a selective destruction of crabs occurs, the broad-fronted crabs being killed in greater proportion than the narrow-fronted crabs.

Therefore, if we could take a number of young crabs, and protect them through a certain period of their growth from the action of this selective mud, the broad-fronted crabs ought to have as good a chance of life as the rest; and in consequence the protected crabs should contain a larger percentage of broad individuals than wild crabs of the same age; and the mean frontal breadth of such a protected population ought to be greater, after a little time, than the mean frontal breadth of wild crabs, in which the broad individuals are being constantly destroyed.

It is difficult to perform this experiment, because one cannot know the age of a crab caught on the shore. But so far as one can judge the age of a crab by its length, I can show you that the thing which ought to happen, on the hypothesis that such selective destruction is going on, does actually happen.

I established an apparatus consisting of some hundreds of

numbered glass bottles, each bottle being provided with a constant supply of clean sea-water by means of a system of glass syphons. Into each of these bottles I placed a crab from the beach. After a considerable number of deaths had occurred, a series of crabs was finally established, each crab living in a numbered bottle, until it had cast its shell. The process of moulting involves no distortion of the carapace, which could affect the measurements concerned, and therefore each cast shell was carefully measured. The measurements of these shells were carefully compared with measurements of wild crabs of the same size, and the mean frontal breadth of these shells was a little less than the mean breadth in wild crabs of corresponding length.¹

After each crab had moulted, it was left in its bottle until it had grown and had hardened a new shell. It was then killed and measured, and the measurements obtained were compared with measurements of wild crabs of corresponding size. This time the captive crabs were unmistakably broader than wild crabs of their own size, and there were a few of the protected crabs which were very remarkably broad. The distribution of abnormalities before and after moulting is shown in Fig. 6.

This is precisely the result which we ought to have obtained, if the hypothesis suggested by the study of mud were true. By protecting crabs through a period of their growth, we ought to raise the mean frontal breadth, and to obtain a greater percentage of abnormally broad crabs, and that is what we have seen to occur.

Of course, this experiment by itself is open to many objections. The estimate of age by size is a dangerous proceeding, and it is difficult to exclude the possibility that confinement in a bottle may directly modify a crab during the critical period of




FIG. 6.—Distribution of abnormality of frontal breadth ratios in 527 female crabs before and after moulting in captivity. The continuous line shows the distribution before, the dotted line after moulting.

moulting, and so on. All these points would have to be discussed at greater length than your patience would bear, before we could accept this experiment by itself as a proof that some selective agent exists on the shore, which is absent from the bottles. At the same time, the result of this experiment is exactly what we should expect to find if such a selective agent did exist, and so it is in complete harmony with the evidence already put before you.

Of course, if the observed change in frontal breadths is really the result of selection, we ought to try to show the process by which this selection is effected.

This process seems to be largely associated with the way in which crabs filter the water entering their gill-chambers. The gills of a crab which has died during an experiment with china clay are covered with fine white mud, which is not found in the gills of the survivors. In at least 50 per cent. of the cases, this difference is very striking; and the same difference is found between the dead and the survivors in experiments with mud.

I think it can be shown that a narrow frontal breadth renders one part of the process of filtration of water more efficient than it is in crabs of greater frontal breadth.

¹ This was probably due to the death-rate during acclimatisation being selective. It was very difficult to keep the apparatus clean; and the deaths which occurred were in most cases due to the presence of putrescent bits of food, which had not been removed.

A subsequent experiment was made with the same apparatus, in which crabs were kept in putrid water until a large percentage had died; and the mean frontal breadth of the survivors was found to be distinctly less than the mean frontal breadth of the dead.

It would take too long to go into that matter now, and I shall not attempt to do so. I will only now ask you to consider one or two conclusions which seem to me to follow from what I have said.

I hope I have convinced you that the law of chance enables one to express easily and simply the frequency of variations among animals; and I hope I have convinced you that the action of natural selection upon such fortuitous variations can be experimentally measured, at least in the only case in which any one has attempted to measure it. I hope I have convinced you that the process of evolution is sometimes so rapid that it can be observed in the space of a very few years.

I would urge upon you in conclusion the necessity of extending as widely as possible this kind of numerical study. The whole difficulty of the theory of Natural Selection is a quantitative difficulty. It is the difficulty of believing that in any given case a small deviation from the mean character will be sufficiently useful or sufficiently harmful to matter. That is a difficulty which can only be got rid of by determining in a number of cases how much a given variation does matter; and I hope I have shown you that such determination is possible, and if it be possible, it is our duty to make it.

We ought to know numerically, in a large number of cases, how much variation is occurring now in animals: we ought to know numerically how much effect that variation has upon the death-rate; and we ought to know numerically how much of such variation is inherited from generation to generation. The labours of Mr. Galton and of Prof. Pearson have given us the means of obtaining this knowledge: and I would urge upon you the necessity of obtaining it.

For numerical knowledge of this kind is the only ultimate test of the theory of Natural Selection, or of any other theory of any natural process whatever.

SECTION G.

MECHANICAL SCIENCE.

OPENING ADDRESS BY SIR JOHN WOLFE BARRY, K.C.B., LL.D., F.R.S., PRESIDENT OF THE SECTION.

APART from all the other considerations which so favourably affect this Congress, I think, so far as Section G is concerned, that we are fortunate in meeting in this ancient city, which has so much of special interest for engineers and for others interested in applied science.

(1) I propose, therefore, to say a few introductory words about Bristol and its neighbourhood from the point of view of this section of the Association, but it is far from my intention to either criticise the past work of the Corporation in relation to their dock enterprises or to volunteer advice to them with respect to possible works of improvement.

Bristol is, at this moment, of great commercial importance, as indicated by the value of its imports and exports, and occupied an even more important relative position among British ports at a time when the ports of Liverpool, Glasgow, Cardiff, or Southampton were almost, or altogether undeveloped. So far as Customs Revenue is concerned Bristol now stands third, and in regard to the gross value of her sea-borne trade she is thirteenth among ports of the United Kingdom.

It is unnecessary, and it would be foreign to the objects of Section G, to attempt to trace the economic reasons which have caused the long-continued importance of Bristol, or to account for the rapid growth of other ports more or less competitive with her. All such causes are to be found, at least to a great extent, in considerations apart from the merely physical characteristics of the sea, river, or land at the various sites, as, for example, in propinquity to markets or centres of production, in situation relatively to population or to means of distribution, in individual or collective enterprise, in enlightened or unenlightened administration.

These circumstances have, in truth, at least as much if not more influence in determining the history and prosperity of ports than what are termed natural advantages of respective sites, by which I mean such matters as protection from winds

and currents, depth of water in the port itself and in its approaches from the sea, the possession of soil adapted to the foundations of docks or quays, and ready access to suitable materials for cheap and efficient construction.

While recognising to the full the great advantages of such physical endowments in the development of a great port, one cannot but remember that they form only part of the problem, and that the business of engineers is to modify and direct the great forces and characteristics of nature for the use and convenience of mankind. We have, in fact, to make the best of a locality which may or may not be promising in the first instance, and history shows us that there are few places which are hopeless for our purposes. Thus while, on the one hand, we see many harbours in this country which inherit from nature every feature to be desired for the establishment of a port, but which remain useless for that object, so, on the other hand, we find many of the great centres of trade established in situations which possessed no such advantages, and where almost everything has had to be supplied by painful exertion and great expenditure.

As examples of these facts, I may point to the remarkable progress of many commercial ports situated in localities which were originally the reverse of promising from an engineering point of view—to Glasgow, where twenty-six millions sterling in value of exports and imports are annually dealt with in ships of the largest draught, though it is placed on a river which only fifty years ago was nearly dry at low water for a distance of ten miles below the present docks—to Newcastle, with a present trade of 13½ millions sterling, which within the memory of this generation was approached by a shallow river, entering a much-exposed part of the North Sea over a dangerous sand bar. Sixty years ago the Tyne could only receive (and that only at high water) a small class of coasting vessels, whereas it is now navigable for deep-draughted vessels for a distance of thirteen miles from the sea. The breakwaters also at Tynemouth, which have been constructed under great difficulties on a coast without a single natural encouraging characteristic, not only make a valuable harbour of refuge, but have, practically speaking, removed the external bar.

In a similar way, as evidence of the truth of my proposition, I might point to a multitude of other instances; to the great docks of Buenos Ayres, which city, when I knew it twenty years ago, could not be approached within seven or eight miles by sea-going ships of fifteen or sixteen feet draught; to Calcutta, dependent on the dangerous navigation of the Hooghly, including the dreaded James and Mary shoals; to the creation of the port of Manchester, forty-five miles from the sea, approached by a tide-locked canal which has cost thirteen or fourteen millions of money in its construction; to the great recent developments of Rouen, Dunkirk, Antwerp, and Amsterdam; to the improvements of the Danube and the Mississippi. In all of these cases the natural characteristics of the localities were quite unsuited to the requirements of an advancing trade in modern vessels, but the inexorable demands of commercial shipping have created the supply, at the hands of engineers, of improvements and modifications of nature, which are so large and important that, to an unprofessional eye, they might now almost appear, at least in some of the cases which I have mentioned, to be physical characteristics of the locality.

I think that we may safely say that trade will produce the required accommodation, and that accommodation in itself will not create or attract trade.

Bristol is a case in point, and it is interesting to us at this meeting to note, however briefly, some of the important works which have altered and are altering its capacity as a port. At the end of last century Bristol and its capabilities were, as they have been almost ever since, the battlefield of civil engineers, and we know that reports and projects were made by most of the men who were then recognised as authorities. The diversion of the river Avon and the construction of the floating harbour of Bristol, which were carried out under the advice of William Jessop in the years from 1804 to 1809, were boldly conceived and ably executed. The result of the diversion of the Avon by means of what is still known as the New Cut enabled the old course of the river to be made into a floating harbour of about 71 acres, of which 57 acres are available for vessels of considerable size. The total cost seems to have been about 600,000*l*. Though the greatest draught of water in the floating harbour (some 20 feet) and the dimensions of the original locks (150 feet long and 36 feet wide) may appear to us at the close of the

nineteenth century somewhat insignificant, they were, no doubt, up to the estimated requirements of that day, and I think we can recognise in Jessop's work the impress of a great mind.

The Cumberland Basin was deepened and improved, and the lock accommodation was increased by Brunel in 1850 by the construction of a lock, 350 feet long and 62 feet wide, and again by Howard in 1871, who made another lock, 350 feet long, 62 feet wide, with 23 feet of water at high water of neap tides. This is the present limitation of the access of shipping to the town docks, and though we realise its insufficiency for modern vessels, we can appreciate the energy of those who have gone before us, and who found the funds for or designed works which have for so many years well fulfilled their purpose.

The approach to Bristol from the sea—that is to say, from King Road in the Bristol Channel—is certainly unpromising for large ships, and indeed, when contemplated at low water, appears not a little forbidding. Something has been done, and more is now in progress, towards straightening, deepening, buoying, and lighting the tortuous course of the Avon below Bristol. More, no doubt, would have been undertaken in former years, if the great rise of tide in the river had not provided, at spring tides, a depth and width for navigation which were sufficient for practical purposes, until the size of modern ships imperatively demanded increased facilities of approach. I think it is a remarkable thing that vessels of 3000 tons burden, 320 feet in length, and drawing 26 feet of water, succeed in reaching Bristol, and that the trade in the heart of the city continues to increase.

Those acquainted with the strong tides of the Avon, or with its bends, which do not exceed in places a radius of 800 feet, and, lastly, with what might be the consequences of a long vessel grounding in a channel which has only a bottom width of 100 feet, cannot but recognise the skill and nerve of the pilots in navigating large vessels from King Road to Bristol. This is done by night as well as by day, and so successfully that the rate of insurance for Bristol is no more than it is for Avonmouth or Portishead, the entrances of which are in the Severn, or than for many ports situated on the open sea.

We have similar examples of what can be done by the systematic development of pilotage skill in the Hooghly, the River Plate, the Yangtse Kiang, the Mississippi, and other rivers where special men have been evolved, as it were, by the demand, and navigate with safety and success channels which are so full of dangers that they might well appear impracticable. Experience, indeed, shows us that, given a trade and a depth of water rendering access possible, ships will make their way to ports through all kinds of difficulties and with a wonderfully small margin of water under their keels, reminding one of the boast of the Mississippi captain that he could take his steamer wherever the channel was a little damp.

To return, however, to Bristol and the Avon; in spite of all efforts to keep pace with trading requirements, the time arrived, in 1868, for providing improved dock accommodation, which would avoid the navigation of the Avon, and at the same time afford deeper locks and more spacious quays than could be given in Bristol itself. The Avonmouth and Portishead docks accordingly were built between 1868 and 1878, and acquired by the Corporation in 1884. Both are fine works for their period; but even in their case the rapid development of modern shipping has occasioned a demand for enlargements of the facilities which they afford. Accordingly, a matter which is again agitating Bristol is still further dock accommodation, and there has been a sharp contention whether this should be effected by what is implied in the somewhat barbarous word "dockising" the Avon, or by new docks at King Road. Dockising implies the construction of a weir and locks at Avonmouth, so that the Avon would be impounded and make one sheet of water nearly six miles long to Bristol, the natural discharge of the river being provided for by outfall sluices, while the alternative of dockising the Avon is to be found in great additions to the docks either at Avonmouth or Portishead.

In the peaceful atmosphere of Section G, I will not enter upon the various aspects of these antagonistic proposals, and will merely say that I have no doubt that in some way Bristol will keep ahead of what is wanted, and that I wish the city and the engineer who may carry out any of the ideas which may be eventually adopted every success and satisfaction in such important undertakings.

(2) Leaving, then, for the present all local considerations, and seeing that a large part of my own work has lain in the

construction of new docks and in the alteration of old docks, I propose now to say a few words on what appear to me to be at present the salient points on these subjects in relation to the growth and the requirements of our merchant navy.

In the first place one cannot but be struck with the great demands which have come with some suddenness on the present generation for increased dock and quay accommodation. The British people are the chief carriers of the world, and are indeed those "that go down to the sea in ships, and occupy their business in great waters." This can be appreciated when we consider that annually our over-sea import registered tonnage is thirty-four millions, and our export registered tonnage is thirty-eight millions. Our coastwise traffic amounts to sixty-three million tons per annum, making together a tonnage to be dealt with of one hundred and thirty-five million tons. If we add to these figures the tonnage of vessels in ballast and the number of calls of those vessels in the coasting trade which touch at several ports in the course of one voyage, we must add a further fifty-five millions of tonnage, making a total of one hundred and ninety millions of tonnage using our ports yearly; and if we divide these figures by, say, three hundred days, to provide against more or less idle days, bad weather, and the like, we have the result of six hundred and thirty-three thousand tons per diem entering and leaving our ports. If we assume an average ship of three hundred registered tons, which is probably not far wrong, we have about two thousand one hundred trading vessels entering or leaving our ports daily—a flotilla of startling numbers.

In truth, the magnitude of our mercantile navy, as compared with that of other countries, is astonishing. We have ten and a half millions of tons, against a total of thirteen millions of tons belonging to all the other nations of the world, in which are included three millions of tons of steam vessels engaged in the lake and river traffic of the United States. Descending to particulars, our merchant fleet is eleven and a half times that of France, seven times that of Germany, eighteen times that of Russia (in Europe), two and three-quarter times that of the United States (inclusive of the craft on the great lakes), six and three-quarter times that of Norway, fourteen times that of Italy, and fourteen times that of Spain. Out of our total tonnage of ten and a half millions, six and three-quarter millions are steam vessels, and the proportions in relation to the steam tonnage of the other countries above referred to are approximately the same.

Again, it is instructive to note how small a proportion of the trade of other countries, even including coasting traffic, is carried in ships belonging to the country in question. Thus, whereas we as a nation convey in steamships 76 per cent. of the aggregate tonnage of our own ports, only the following proportions of the total trade of other nations are carried by the shipping of each country in question:—

France	about 30 per cent.
Italy	19 "
Germany	43 "
Russia (in Europe)	7 "
Norway	56 "
Sweden	29 "
Holland	26 "
United States (over-sea)	15 "

Further, it is a recognised fact that a very large part of the balance of the above proportions is conveyed in British ships frequenting the various foreign ports and acting, as I have said, as the ocean carriers of the world.

Thus in the best returns available I find that British shipping conveys the following proportions of the over-sea commerce of other countries:—

Italy	44 per cent.
Germany	38 "
Russia	57 "
Norway	18 "
Sweden	27 "
Holland	54 "
United States	60 "
France	(not given)

The experience of the Suez Canal again tells the same tale, for of the total tonnage passing through that international waterway 66 per cent. is British. This is nearly seven times that of the shipping of the next largest contributor, which is Germany, and nine times that of France.

NO. 1508, VOL. 58]

This vast amount of carrying trade is in British hands, because we can do it cheaply as well as efficiently. I believe that the whole of our commercial fleet is worked at a very narrow margin of average profit, though in the aggregate it forms one of the most important factors in our country's position among the nations of the world.

We are often reminded of how greatly the value of our imports exceeds that of our exports; but we should not forget that the profit on the transport of both goes chiefly to the British nation as shipowners, in addition to the profit which is earned by them in the carriage of merchandise from one foreign port to another.

What an important thing it thus is to the prosperity of this country, not merely that our own ports should be convenient and adequate to all demands, but that our ship-builders should be able to keep pace with the demands of this huge transport traffic! We find in this connection that we add about half a million of tons of shipping annually to our register, and that we lose about 250,000 tons annually by wreck and by vessels becoming old or obsolete, so that, as a matter of fact, the average annual increment of our mercantile navy for the past twelve years is about a quarter of a million of tons.

The remarkable development within recent years in the cheapness of steam navigation, the improved methods of building and rigging of sailing ships, and various economic causes have resulted in a large increase of the average size of ship engaged in over-sea voyages with a comparative diminution in the number of the crews of each description of vessel. Greater draught of water is consequently demanded, and as a better knowledge of ship-building has indicated that the beam of ships can be considerably increased without involving greater resistances, we may expect to see ships to increase not only in length and depth, but also in width.

The largest steamer twenty years ago (excepting of course the *Great Eastern*, which was a magnificent conception, though in advance of her time and its requirements) was, I believe, the *City of Berlin*, of 5500 tons burden. Her length was 488 feet, and her draught and beam were 25 feet and 44 feet respectively. At the present time the *Kaiser Wilhelm der Grosse* is 625 feet long, her beam is 66 feet, and her draught is 27 feet, and we know that these dimensions will soon be exceeded.

A modern liner now being built will have a length of 704 feet (or 24 feet longer than the *Great Eastern*) with a beam of 68 feet and a draught of 28½ feet. The great steamers for the transport of cattle are 585 feet long, 64 feet beam, and 30 feet draught and upwards, carrying 14,000 tons of cargo. Some of the large sailing vessels carry over 6000 tons dead weight and draw 28½ feet. Ships of war, though not so long as liners, have a beam of 75 feet with a draught of 31 feet, and though in the commercial marine we need not perhaps anticipate any great further increase of draught of water, the demand for which is largely governed by what is available in foreign ports or rivers and in the Suez canal, the fact that men-of-war can, with due regard to economy of propulsion, be built with great width of beam in proportion to length, seems to indicate that we must be prepared in the future for a considerable increase of beam for cargo-carrying vessels.

We have further to note that, owing, no doubt, to the vast improvements of marine steam engines and boilers realising unlooked-for economy in the combustion of coal, steam vessels are supplanting all but the largest class of sailing vessels as carriers of commerce, almost as rapidly as they did forty or fifty years ago in the conveyance of passengers and as ships of war.

In 1897, out of a total shipping trade (cargoes and ballast) dealt with in ships of all nations at the ports of the United Kingdom, amounting to ninety millions of tons, eighty-one millions of tons, or 90 per cent., were conveyed by steam vessels; whereas, in 1885, out of a total of sixty-four millions of tons, fifty millions of tons, or 78 per cent., were in steamers. If we take, however, the tonnage of cargoes and ballast conveyed to and from her own ports by British ships only, we find that in 1897, out of a total of sixty-four millions of tons, sixty-one millions of tons, or 95 per cent., were in steam vessels; whereas, in 1885, but 85 per cent. of the total tonnage conveyed by British vessels was in steamships.

Of the tonnage of vessels built in the United Kingdom in 1885, 50 per cent. were steamers, but in 1897 the proportion was 86 per cent.; and to sum up, we find that in the commercial fleet of the United Kingdom and British Possessions, as between 1887 and 1897, sailing ships have decreased 16 per

cent. in number and have, in spite of the building of a certain number of exceptionally large vessels, decreased 9 per cent. in average size; while steamers have increased 23 per cent. in number and 16 per cent. in average size. The total sailing tonnage has decreased in the same period by 24 per cent., and the steam tonnage has increased by 36 per cent.

The problems thus confronting us, as results of the increased size of all descriptions of over-sea steamships, require much consideration from an engineering point of view, and are further puzzling, and will continue to puzzle, our financial authorities, without whose aid the engineer can do but little.

We ask, Where is all this expansion of requirements to stop, and how far are we justified in extending our view of the wants of the future from the contemplation of the conditions of the present and of what has occurred in the past? This is undoubtedly a difficult question, and he would be a bold man who thought that we had reached finality in the size of ships. Bound up with this consideration are not merely matters of first cost of the accommodation to be provided, but also of the annual expenses in working and maintenance, not only of the docks themselves, but in what is perhaps of more importance, viz. the preservation of sufficiently deep and wide approaches to them.

Apart from length, depth, and beam, the midship cross section of modern cargo ships has altered completely of late years, and is now nearly as rectangular in shape as a packing-case, excepting only that at the bilges the sides and floor are joined by a curve of small radius. The keel has almost disappeared, and bilge keels are often added. The result of these alterations of shape in the ordinary hulls of trading ships is that the sills and sides of many locks and entrances are now unsuited to what is wanted, and consequently their original power of accommodating vessels is most seriously diminished.

Until lately it was generally considered that locks 600 feet long, 80 feet wide, and 26 feet deep were sufficiently capacious, with some margin for future wants; but I think we must now go further in length and depth, and not improbably to some extent in width. We find that at Liverpool the Dock Board have ordered vestibule basins to act as locks 1150 feet long and 520 feet wide, with entrances 100 feet wide and 32 feet deep; and somewhat similar dimensions were talked of for the entrance lock of the recently proposed Windsor Dock at Penarth, which was intended to be 1000 feet long, 100 feet wide, and 34 feet deep at neap tides.

Again, apart from the question of locks and entrances, the older docks themselves are beginning to be found too shallow and too narrow for modern vessels. In docks which are deep enough at spring tides and too shallow at neap tides, and which are opened to the "tide of the day," much may be done to improve the depth by systematic pumping, so as to keep the surface always at the level of high water of spring tides. By this expedient, large areas of old docks may be to that extent modernised at the expense, perhaps, of new entrance locks and the annual cost of pumping. This latter yearly outgoing is not an important matter. At Liverpool and Birkenhead 230 acres of nearly obsolete docks have been thus improved at a capital cost of about 96,000*l.* for pumping machinery and an annual expenditure of 6000*l.* I am executing a similar improvement by pumping in one of the smaller docks on the Thames, and contemplate it on a larger scale at an important dock there, and also at Hull.

The conditions of commerce now require also, in order to realise the necessary economy of transport, the greatest despatch, for demurrage on the large and expensive modern steam vessels is a most serious question. Thus there must now be no waiting for spring tides, or, if possible, for rise of tide on the day of arrival. Every steamer expects to discharge her cargo on to the quay without waiting for much stacking, still less for trucks; and under modern conditions dock work must be got through in one-third of the time which was considered proper ten or twelve years ago. From these reasons larger quays and warehouses, better railway approaches, improved sidings, and better machinery are all necessities, as well as deeper water and better approaches.

These demands have come on us, as I have said, not so much gradually as more or less suddenly, and the call for improved docks is general, and, in my opinion, it will be continuing.

Liverpool last year undertook to spend nearly five millions on such works, and we know of very many important projects at other places. Taking the expenditure within the past decade,

and adding to it the authorised expenditure at Liverpool, at the great ports on the Bristol Channel, on the Thames, at Southampton, Hull, Middlesbrough, Hartlepool, Sunderland, the Tyne and its neighbourhood, at Grangemouth, the Fife Ports, at Glasgow, the Ayrshire Ports, the Cumberland and Lancashire Ports, and so round the British coasts to Preston, I roughly estimate an expenditure, either made during the past ten years or contemplated, of from 35 to 40 millions.

These are large figures, and we ask from whence will an adequate revenue come; for it is a more or less accepted fact that docks by themselves do not produce more than a very moderate return on their cost, though, of course, there may be exceptions to every rule. Apart from the expenditure which has been undertaken much remains to be done, and the source of supply of the capital required is a highly important consideration. I venture to think on this point that we should learn to realise that under modern conditions docks should be considered largely in the light of being railway stations for goods and minerals and, in many cases, for passenger traffic. Docks and quays, together with improved approaches from the sea, are, in fact, the means of bringing traffic to the railways (and, to a less degree, to the canals) of a country, and should be looked upon as links in the chain of transport and inter-communication.

They are certainly as necessary adjuncts of a railway, at least in our country and in respect of goods and minerals, as large stations and depôts are in all important towns.

The older view of our Parliament was that docks and railways should be in different hands; but I much question whether this idea should now commend itself. It is difficult, as I have said, for a dock enterprise standing alone to make any considerable return on its cost, and though it is true that capital can be found under guarantees of an already developed trade by some of the great Dock Trusts, such as at Liverpool or Glasgow, the return is but a modest one, and not such as is likely to tempt capitalists to new ventures in constructing or enlarging many of the docks which stand in need of improvements.

On the other hand, a railway company which gets a fairly long lead for the goods to and from a dock can afford to look at the matter of expenditure on docks with some liberality. We have conspicuous examples of great public benefit being afforded at Southampton and at Hull, where the docks have lately passed from the hands of financially weak companies dependent only on dock dues, to the ownership of powerful railway companies. Similarly, several of the north-eastern ports besides Hull—the large docks at Grangemouth, Barry, Penarth, Garston, Fleetwood, and elsewhere—are further examples, amongst others, in which the revenue of railway companies has been spent on dock improvements with a spirit which would be otherwise unattainable. A dock also must necessarily be nowadays almost wholly dependent for its efficient working on the best understanding being maintained with the railway companies for the prompt and adequate provision of land transport, so that in that point of view also the two interests are one and should be recognised as such.

In the consideration of the advisability for concentration of ownership, there remain only the questions of safeguards against unfair treatment of competitive modes of transport, such as canal and road traffic, and provision against any improper results of monopoly of railway access. These, I think, can be provided by Parliamentary enactment, either by insisting on adequate access under proper conditions for all within reach, or, in any case, of inadequate facilities being accorded, by authorising the construction of other docks in the hands of competing railway companies or of other aggrieved parties, with in such cases railway privileges. With these safeguards the public could be efficiently protected, and, if this be so, I cannot but think that, *ceteris paribus*, the trading community will be better served by docks directly connected with railway companies than by separate existences and management. On the one hand, I hope that those who administer the great railway undertakings will realise this community of interest, and, on the other, that Parliament will favour intimate financial relations between docks and railways, instead of more or less systematically discouraging such connection. This question is one which is peculiarly interesting here at Bristol, where the docks are in the hands of the Corporation, and where the railway companies carry the traffic, which, but for the docks, would be largely non-existent.

(3) Leaving now the question of modern docks and shipping,

as to which, as I have said, Bristol is interesting to engineers, there are one or two other matters of history which appeal to Section G in this locality. In the first place, Bristol was the birthplace of the Great Western Railway. I. K. Brunel, its engineer, had previously, by public competition, been selected to span the gorge at Clifton by a suspension bridge of the then almost unrivalled span of 702 feet. Again, under the influence of Brunel, Bristol became the home of the pioneers of Transatlantic steamships, and the story of the initiation of the enterprise is thus told in the memoirs of his life. In 1835, at a small convivial meeting of some of the promoters of the Great Western Railway, some one said, "Our railway to Bristol will be one of the longest in England," and Brunel exclaimed, "Why not make it the longest line of communication in the world by connecting it with New York by a line of steamers?" Out of this grew the *Great Western* steamship, and the history of the enterprise and of its success is too well known, at least here, to require any allusion to the steps by which it was brought about. Suffice it to say that, in spite of much discouragement, the *Great Western*—of the then unexampled size of two thousand three hundred gross tons, and with engines of unparalleled power—was launched at Bristol in 1837, and ran successful and regular voyages till 1857, when she was broken up.

In Section G there are many who can appreciate the difficulties of such a new departure as the *Great Western* steamship, even if they had been confined to the design and study of a vessel and engines of unprecedented size; but it is not easy to realise the anxiety and trouble caused by the dictum of a man of science so universally admired as Dr. Lardner, at the meeting of the British Association in this city in 1836, that the whole idea of ocean navigation on voyages as long as from Bristol to New York was at that epoch an abstract impossibility.

In these days of criticism of the past, often involving the rehabilitation of individuals, it is interesting to note that Dr. Lardner's part in condemning beforehand the construction of the *Great Western* steamship and the ideas on which she was designed has been of late years unduly minimised. It has been said that all Dr. Lardner meant was to express a pious doubt as to the commercial prospects of ocean navigation. I have carefully read the *Proceedings* of the time, and I am brought to the conclusion that his words and writings will admit of no such interpretation. Dr. Lardner's views, arrived at after calculation and reasoning, were precisely expressed and boldly and honestly enunciated by him. The words of the discussion here appear not to have been preserved, but in an elaborate article in a *Quarterly Review* in 1837, which is, I believe, admitted as having been written by Dr. Lardner, he said, "that in proportion as the capacity of a vessel is increased, in the same ratio, or nearly so, must the mechanical power of the engines be enlarged and the consumption of coal augmented." He based his views that success was impossible on principles which he supposed to be sound, but which were, in fact, assumptions—viz. that the resistance to the progress of a ship varied directly with her capacity, that a certain number of tons of coal were required per horse-power for the voyage across the Atlantic, and that, this being so, enough fuel could not be carried in a ship, however large she might be made.

Brunel, on the other hand, contended that Dr. Lardner's views were fundamentally erroneous; for that, whereas the capacity of a ship increased in the ratio of the cube of her dimensions, the resistance to her progress varied more nearly as the square. Thus, by adopting a proper length, beam, and draught, a ship would not only carry coal for the journey to New York, but be commercially successful in respect of cargo and passengers.

It is interesting to note that 9 lbs. of coal per indicated horse-power per hour (as compared with our present 1½ to 2 lbs.) was the approximate coal consumption which was more or less accepted by both sides in the controversies of 1836 and 1837.

We know now that the resistances encountered by a ship are not merely dependent on her dimensions, but comprise wave-making at various speeds, bringing form and proportion of dimensions largely into the necessary calculations; but I want to point out that the line of divergence of the different views of Lardner and Brunel was sufficiently precise and quite crucial. It is true that Dr. Lardner, in later criticisms of 1837, retreated somewhat from his position of 1836, introducing more of the commercial aspect of the case and stating that no steam vessel could make profitable voyages across the Atlantic, at

least until marine engines were immensely improved; but, even so, it seems clear that the fundamental matter at issue in 1836 and 1837, the period of Dr. Lardner's active criticism, was the question of the resistances increasing in the same ratio as the capacity. The results of these *ex cathedra* statements by Dr. Lardner about the *Great Western*, then in process of being built, must have caused great anxiety to the promoters and much preliminary distrust of the ship on the part of the public. They were, unquestionably, honestly arrived at, however much they were due to reasoning on unascertained premises, and this latter is the reason for my venturing now to refer once more to them. As a matter of fact, the ship started from Bristol in 1838, and arrived at New York in fourteen days with 200 tons of coal in her bunkers.

Let me remind you of another somewhat similar instance of the way in which the anxieties of engineers have been unnecessarily increased and public alarm gratuitously, though honestly, aroused. When the designs of the Forth Bridge—of which the nation, and indeed the world, is proud—had been adopted both by the Railway Companies who were to find the capital and by Parliament, a most distinguished man of science—the then Astronomer Royal—came to the conclusion that the engineers had neglected certain laws which he enunciated respecting the resisting power of long struts to buckling, and that the bridge ought not to be constructed, as he considered that, to use his own words, "we may reasonably expect the destruction of the Forth Bridge in a lighter gale than that which destroyed the Tay Bridge." All this was stated no doubt: from a strong view of public duty, in a letter to a public newspaper, though subsequently and frankly withdrawn. If the bases of his calculations were right, the conclusion might have been correct; but the fact was, that there was no foundation worthy of the name for the reasoning. Again, another distinguished mathematician publicly criticised the Forth Bridge with equal vigour, basing his views that it was fundamentally incorrect on another set of equally erroneous assumptions, maintaining again that it should not be permitted, because he proved by reasoning on those assumptions that it must be absolutely unsafe.

Once more, in ship-building, until Mr. William Froude, some years prior to 1875, made his experiments by means of models on the highly difficult and otherwise almost insoluble causes of the retardation of ships and their behaviour in waves, beginning at the beginning, taking nothing for granted, and eliminating all elements of possible errors, little or nothing was known of the laws governing these questions. Laws had been laid down by high authorities as to the causes of retardation of ships, many of which, in fact, were not true, while some of the assigned causes were non-existent and some real causes were unrecognised. Mr. Froude was told that no information could be learnt from experiments on models which would be applicable to full-sized ships, and that ships must continue to be designed and engines built on data which, scientifically speaking, were assumptions. The outcome has been that Mr. Froude's *a priori* depreciated experiments with models have solved most of the questions relating to that branch of naval architecture; and at the present time every ship in the Royal Navy, and not a few in the merchant service, are designed in accordance with the data so gained.

Another example of hasty generalisation occurs to me, and that is on the important question of wind pressure. Tredgold, who undoubtedly was one of the soundest of engineers, laid down in 1840 that a pressure of 40 lbs. per square foot should be provided for; reasoning, no doubt, from the fact that such a pressure had in this country been registered on a wind gauge of a square foot or less in area. As a consequence, he assumed that the same force could be exerted by the wind on areas of any dimensions. Thus roofs and bridges, wherever any calculations of wind pressure were, in fact, made, were designed for a pressure of 40 lbs. per square foot of the whole exposed surface, and under the alarm caused by the fall of the Tay Bridge in 1879, the piers of which were not probably strong enough to resist a horizontal pressure of one-fifth of such an amount, a further general assumption was made, and railway bridges throughout the kingdom were ordered by the Board of Trade in 1880, acting no doubt on expert advice, to be in future designed, and are designed to this day, to resist 56 lbs. of horizontal wind pressure on the whole exposed area with the ordinary factors of safety for the materials employed, as if such horizontal strain were a working load.

It had, for a long time previously to this order of Government being issued, been suspected that these small-gauge experiments were untrustworthy, and subsequent experiments at the Forth

Bridge on two wind gauges of 300 square feet and of $1\frac{1}{2}$ square feet respectively, indicated that with an increase of area the unit of pressure fell off in a very marked degree. Under the same conditions of wind and exposure, the larger gauge registered a pressure 38·7 per cent. less per square foot than the smaller gauge. I have been able to carry experiments further at the Tower Bridge by observing the pressure on the surface of the bascules of the bridge as evidenced by the power exerted by the actuating engines. In this case we have a wind gauge of some 5000 feet in area, and it has been shown that, while small anemometers placed on the fixed parts of the bridge adjoining the bascules register from 6 to 9 lbs. per square foot, the wind pressure on the bascules is only from 1 to $1\frac{1}{2}$ lbs. per square foot.

It is difficult to imagine the amount of money which has been expended in unnecessary provision against wind strains of 56 lbs. per square foot on large areas in consequence of this hurried generalisation from insufficient data. I know something of what the provision for 56 lbs. on the square foot for wind cost at the Tower Bridge, and I do not wish to mention it; but if the public had been told that the dictum of experts, arrived at however hastily in 1880, was to be set aside in the construction of that bridge, all confidence would have been beforehand destroyed in it, and I suppose no Committee of Parliament would have passed the Act.

I have mentioned these matters, which could be added to by many similar instances in other branches of applied science, not for the sake of reviving old controversies or of throwing a stone at highly distinguished men, honoured in their lifetime and honoured in their memory, nor for the sake of criticising more modern men of science or a Government Department. Still less do I wish to question the necessity and value of mathematical calculations as applied to the daily work of engineering science, but I recall the circumstances for the purpose of once more pointing out the extreme value of experimental research and of bespeaking the utmost caution against our being tempted to lay down laws based on unascertained data. We know the tendency there has been at all times to generalise and to seek refuge in formulae, and we cannot but know that it is not at an end now. We ought to recognise and remember how few physical questions had been exhaustively examined sixty years ago, and may I say how comparatively few have even now been fundamentally dealt with by experiment under true scientific conditions? The investigation of physical facts under all the various conditions which confront an engineer requires much care, intelligence, time, and last, not least, not a little money. In urging the vital necessity of investigations, I am sure that I shall not be understood as decrying the value of the exact analysis of mathematics, but we must be quite sure that the premises are right before we set to work to reason upon them. We should, then, exert all our influence against rules or calculations based merely on hypothesis, and not be content with assumptions when facts can be ascertained, even if such ascertainment be laborious and costly. In a word, let us follow sound inductive science, as distinguished from generalisations; for "Great is truth and mighty above all things."

In connection with this subject, I may congratulate the Association generally, and this Section in particular, that there is now more hope for experimental science and some endowment of research in this country than at any former time. The vital necessity of further work in these directions has long been recognised by men of science and was notably urged by Prof. Oliver Lodge. Last year, in no small degree owing to the exertions of Sir Douglas Galton, K.C.B., who presided over the British Association in 1895, and brought the question very prominently forward in his inaugural address on that occasion, a highly influential deputation waited on the Premier to urge that England should have a Public Physical Laboratory at which facts could be arrived at, constants determined, and instruments standardised. The importance of the questions which could be determined at such an institution in their influence on the trade and prosperity of the country, independently of the advancement of purely scientific knowledge, cannot well be exaggerated.

Our Government, while somewhat limiting the scope of the inquiry, appointed a small Committee to examine and report on this highly important subject. It is no breach of confidence to say that the Committee, after taking much evidence, visiting a similar and highly successful institution on the continent, and studying the question in all its bearings, were convinced of the great public benefits which may be expected from such an

institution, and have unanimously reported in favour of its establishment.

I feel sure that we shall all earnestly hope that Government will carry out the views of the Committee, and I venture to suggest that each of us should use what influence he may have, to induce the Chancellor of the Exchequer to find adequate funds for an institution which may be of the greatest benefit not merely to scientific research, but to the commerce of these islands, threatened as it is on all sides by foreign competition of the most vigorous description—a competition which is supported by every weapon which the science of other lands can forge for use in the struggle. It being acknowledged that our own work in life is to deal with physical facts and apply them for the use of our fellow-men, we may have good hopes that at such an institution as I have indicated, directed, as it no doubt will be, by the highest scientific superintendence, we shall be able, at least far better than at present, to have a sound knowledge of many facts which are obscure, and to deal with the many new conditions under which the applied science of the future will have to be carried on.

Those who know most of the problems of nature feel the more strongly how much remains which is unknown and realise how completely those who teach require throughout their lives to be always learners. Let each of us then in our special walk of life, seeking for further enlightenment on the various problems of our work and in the application of that science which we love, humbly recognise that,

"All nature is but art, unknown to thee;
All chance, direction which thou canst not see;
All discord, harmony not understood."

INTERNATIONAL SEA FISHERIES CONGRESS AT DIEPPE.

THE movement for the international discussion of matters connected with the sea-fishing industry has made such progress during the past few years that a summary of the proceedings of the recent international congress held at Dieppe should interest readers of NATURE, especially as the regulation of the industry tends more and more to be determined in accordance with the evidence accumulated by scientific investigators. The Dieppe Congress was organised by the Société d'Enseignement professionnel et technique des Pêches Maritimes, and is the second international congress promoted by that society. The previous congress was held at Sables-d'Olonne in 1896, on which occasion Mr. (now Sir) John Murray was the British representative. More than 300 delegates assembled at Dieppe, among whom may be mentioned Mr. C. E. Fryer, of the Board of Trade; Dr. J. H. Fullerton, formerly of the Scottish Fishery Board; Mr. Walter Garstang, representing the Marine Biological Association; Mr. O. T. Olsen, of Grimsby; Mr. Johnsen, of Hull; Drs. Brunchorst and Bull, of Bergen; Dr. Malm, of Gothenburg; M. Tabary, of Ostend; Prof. Vinciguerra, of Rome; Dr. Valle, of Trieste; Dr. Kishinouye, of Japan; Mr. Thorndike Nourse, of the United States; and of course a large number of French delegates representing the Government and various fishery societies and schools, fishing centres and municipalities, including M. Roché, Inspector-General of Fisheries; Prof. Perrier, Baron Jules de Guerne, MM. Lavieville, of Dieppe; Canu, of Boulogne; Odin, of Sables-d'Olonne; Gourret, of Marseilles; and Le Seigneur, of Granville. The proceedings of the Congress opened on the morning of September 2 with an address from the President, Prof. Ed. Perrier, Membre de l'Institut de France. The greater part of the President's address was devoted to an examination of purely French problems—the relative scarcity of steam trawlers and liners, the need of greater solidarity, of a spirit of co-operation and compromise among rival fishing industries, the present unsatisfactory arrangements—or lack of arrangements—for fishery research. This, he said, seemed to demand the creation of a central Fishery Board for France, similar to that of Scotland, which should be charged with the duty of coordinating the work of the numerous marine laboratories in which fishery research is now carried on without concerted aim. Proceeding then to matters of more general interest, he pointed out the advantages which would ensue if the study of plankton could be put upon an international basis by a regular organisation of the marine laboratories of different countries, or by international co-operation in deep-sea expeditions for the solution of problems

connected with the migrations of fishes. At the same time, he said, it would not do to be too ambitious. The extravagant expectations which were held some years ago as to the beneficial effects of sea-fish hatcheries had not been realised either in America, in Scotland or in Norway. Even if the idea were sound, the actual plan of operation needed modification, since the young fish were being turned into the sea at too early an age. Moreover, he asked, would it not be simpler, and in the end more profitable, to complete the investigation of the whole life-history of valuable fishes before launching upon costly and problematic schemes of fish multiplication? To ensure the adequate discussion of these and similar problems, the President, in conclusion, expressed the intention of himself and his colleagues to propose the creation of a permanent international committee for the organisation of future congresses on sea-fisheries, which would extend and complete the work initiated by the French Society.

The subsequent discussions of the congress took place partly at general meetings, partly at the meetings of different sections. Four of the latter were constituted, viz. (1) Scientific Researches; (2) Fishery Apparatus, Preparation and Transport; (3) Technical Education, and (4) Fishery Regulations. The subjects which came before the general meetings dealt with oyster and mussel culture, provident institutions (insurance against accidents, &c.), international regulations for preventing collisions at sea, and co-operation amongst fishermen. As regards the sectional meetings, the topics of general interest naturally fell chiefly within the scope of the first and fourth sections. In the first section the following were the more important papers read: (1) On the natural history and fishing grounds of the Tunny in the Gulf of Gascony, by M. Odin, in which the author showed that the migrations of the Tunny of these waters are less extensive than was formerly imagined, since the fish can be taken in the Gulf throughout the year, although the actual fishing grounds shift with the seasons; (2) On the natural history of the mackerel, by Mr. W. Garstang, in which it was maintained that, as a result of researches recently carried out by the Marine Biological Association, the common species of mackerel can be subdivided into several local races, viz. an American, an Irish, and a race common to the English Channel and North Sea. These researches lead to the conclusion that the winter haunts of the mackerel cannot be situated far from the localities first visited by the several races in the spring; (3) On a proposed biological and physical investigation of the English Channel during 1899, by Mr. Garstang, in which the author invited the co-operation of French societies and naturalists with the Marine Biological Association for a joint periodic survey of the Channel during the coming year. The proposal was supported by Baron Jules de Guerne and M. Odin, and a resolution on the subject was unanimously adopted; (4) On the sea-fish hatchery at Flodevigen, by Captain Dannevig (read in his absence by Baron de Guerne). This paper gave rise to a vigorous discussion on the efficacy of hatcheries. Captain Dannevig contended that the success of his methods was attested by the statistics of cod taken in Christiania Fjord, but this statement was categorically denied by Dr. Brunchorst, and also adversely criticised by Dr. Fullerton and M. Canu.

The principal papers read in the fourth section were as follows: (1) On trawling in territorial waters, by M. Sauton; (2) On the necessity of new regulations concerning the mesh of drift and fixed nets, by M. Maraud; (3) Trawling and its effects, by M. Coutant; (4) On the territorial limits, by Mr. Olsen. The discussion which followed these papers was long and interesting, but cannot be fully summarised here. It will suffice, however, to say that at the subsequent general meeting of the congress a resolution against trawling (of all kinds) within three miles from low-water mark was carried by 37 votes to 9; and that other resolutions were carried which would have for effect the prohibition of certain kinds of fishing beyond the present territorial limits, and would prohibit the sale of immature fish, the *minimum* size for each species to be fixed hereafter by an international commission of fishermen, owners, public officials, and scientific experts.

It may be stated in conclusion that the memoirs read before the general meetings of the congress are already published (Paris, Augustin Challamel, Rue Jacob 17), and that the papers communicated to the different sections, with the final resolutions of the congress, will be published in a second volume in the course of the next few months.

NOTES.

WE are reminded that the new laboratories of Physiology and Pathology, which University College, Liverpool, owes to the generosity of the Rev. S. A. Thompson Yates, will be opened on October 8 by Lord Lister, President of the Royal Society. By his benefaction, Mr. Thompson Yates has strengthened the medical school of the College in a very marked degree, and has enabled the professors of physiology and pathology to take advantage of the most recent additions to our knowledge in their lectures and laboratory instruction. Lord Lister will be accompanied on the occasion by a large and distinguished party. The Lord Mayor will represent the city; Earl Spencer, Chancellor of the Victoria University, has promised to attend and admit Lord Lister to the degree of D.Sc. conferred on him by the Victoria University; Lord Derby, President of the College, will be present, with the authorities of the Victoria University and its Colleges. Among those who have accepted the invitation of the College Council may be mentioned: the Duke of Devonshire, Lord Derby, Lord Spencer, Lord Ripon, Lord Kelvin, Mr. A. J. Balfour, Prof. Michael Foster and Prof. Rücker (the Secretaries of the Royal Society), Prof. Virchow, Sir Douglas Galton, Sir Samuel Wilks, Sir Richard Thorne, the Bishops of Liverpool, Chester, Carlisle, and Ripon, Sir William Gairdner, Mr. Justice Kennedy, Sir James Crichton Browne, Dr. Lauder Brunton, Sir Archibald Geikie, Captain Abney, C.B., Sir George King, Mr. Thielton-Dyer, Prof. Ramsay, Prof. David Ferrier, Dr. Pavy, Mr. R. B. Haldane, Sir John Batty Tuke, Sir Henry Littlejohn, Prof. Schäfer, Prof. Burdon-Sanderson, Prof. Kanthack, Prof. Halliburton, Prof. Meldola, Prof. Poulton, the Dean of Lichfield, Prof. Charlton Bastian, the Hon. Sydney Holland, Prof. Rose Bradford, Prof. Forsyth, Prof. Bower, Dr. Alexander Cope, Prof. Crookshank, Prof. Waller, Prof. Noël Paton, Dr. Ludwig Mond, Dr. Mott, Prof. Stirling, Prof. Liveing, Mr. Gerald Yeo, Prof. Macalumn, and Dr. Byrom Bramwell. The proceedings will commence with the degree ceremony, which will take place in St. George's Hall at 3 o'clock. Lord Lister will then, with the President, Earl Derby, proceed to open the new laboratories. In the evening a banquet will be given by the Lord Mayor in the City Hall.

ON Sunday, the 11th inst., one of the most destructive hurricanes that has occurred for many years visited Barbados and the Windward Islands, causing immense damage to property and great loss of life. These storms usually occur between July and October, when the equatorial calms are furthest north of the equator; the late A. Poey, of Havana, compiled a list of all hurricanes observed in the West Indies since 1493, and of these nearly 80 per cent. occurred in those months. They usually commence to the eastward, and travel in a north-westerly direction till they reach about latitude 25° N., when they recurve in a north-easterly direction. So far as is known from the meagre reports which have yet been received, this disastrous storm followed the usual track, as the observer of the United States Weather Bureau at Jamaica seems to have forwarded notice through New York that a storm was approaching Barbados from the southward on Saturday, the 10th inst.; but, owing to an unfortunate interruption in the cable, the warning arrived too late. The late Rev. B. Viñes, S.J., formerly director of Havana Observatory, made a special study of West Indian hurricanes during a period extending over twenty-three years, and shortly before his death (in 1893) prepared a paper upon the subject for the Meteorological Congress at Chicago, which is regarded as the most satisfactory statement of the behaviour of these storms that has yet been made. This paper has just been published in a separate form by the United States Weather Bureau. In it the author discusses very completely the general laws of cyclonic circulation and translation,

including the law of the recurving of the path of the hurricanes in the different months of the cyclonic season.

THE Press Association states that on Thursday last Mr. Stanley Spencer and Dr. Berson ascended from the Crystal Palace in a balloon inflated with pure hydrogen gas, and attained the remarkable altitude of 27,500 feet, or only 1500 feet less than Coxwell and Glaisher's highest in 1862. Numerous scientific instruments, including a self-recording aneroid barometer, were carried, and also compressed oxygen for inhaling at the greatest height. The descent was near Romford. At 25,000 feet the air became so rarefied that both explorers had to breathe the compressed oxygen taken with them. The balloon had a capacity of 56,500 cubic feet.

IT is reported through Reuter's Agency that a stream of lava from Vesuvius has destroyed a part of the roadway leading from the observatory to the lower station of the funicular railway. A mass of molten rock is flowing down the mountain side in three streams—one along the foot of Monte Somma, a second through the middle of the Vetrana zone, and a third along Monte Crocella. The stream running round the base of Monte Somma continues to burn the chestnut woods, and nearly reaches the observatory. The central flow has reached a point close to the Carabinieri barracks, while the Crocella stream, after passing close to Messrs. Cook and Son's building, has reached the northern edge of the Canteroni ridge, whence it may also threaten the observatory. News from Naples on Saturday states that the eruption is becoming hourly more active and more menacing, and the streams of molten lava are spreading in every direction. The most threatening is that which is flowing down the immense valley of Vedrino, which is now almost filled. The observatory, which was originally situated at a height of 610 metres, is now said to have sunk over 27 metres owing to the sinking of the ground. Seven new craters have formed round the central crater, without, however, in any way diminishing the activity of the latter. The gravity with which the outbreak is regarded is chiefly based on the fact that the volcano is throwing out stones and scorie similar to those ejected in the great eruption of April 1872, when the lava streams covered an area of two square miles, averaging 13 feet in depth, and the damage to property exceeded three million francs.

THE Berlin Academy of Science has made the following grants for botanical work:—2000 marks to Prof. Eichler, for the continuation of his work on East African plants; 600 marks to Prof. Graebner for the continuation of his work on German heaths; 500 marks to Dr. Loesner, for the completion of his monograph of the Aquifoliaceæ.

TWO Walker prizes, of the value of sixty dollars and fifty dollars respectively, are annually offered by the Boston Society of Natural History for the best memoirs written in the English language on subjects proposed by a committee appointed by the Council. The subjects for 1899 are: (1) Is there fundamental difference between "equation division" and "reduction division" in the division of cells? (2) The phenomena and laws of hybridisation. The subjects for 1900 are: (1) Stratigraphy and correlation of the sedimentary formations of any part of New England. (2) A study in paleozoic stratigraphy and correlation. Memoirs must be sent in on or before April 1 of the year for which the prize is offered.

THE Mayor of Angers has appointed M. Albert Gaillard curator of the Lloyd herbarium in that town.

WE learn, from the *Botanical Gazette*, that Dr. A. Möller, of Eberswald, has undertaken the preparation of a memoir of Fritz Müller, so well known in connection with the Flora of Brazil, and with problems connected with the pollination of plants.

THE death is announced of Dr. H. Trimble, professor of practical chemistry, Philadelphia College of Pharmacy, and editor of the *American Journal of Pharmacy*.

NEWS has been received that Mr. de Windt, the geologist with the Belgian scientific expedition for the exploration of the Congo, was drowned on Lake Tanganyika on August 9, with Mr. Kaisley, a gold prospector.

THE *Pall Mall Gazette* announces that Mr. S. A. Rosenthal and Dr. S. J. von Komocki have succeeded in preparing matches which do not contain yellow phosphorus, and are capable of ignition by friction upon any surface. It is claimed that these matches can be manufactured as cheaply as the ordinary ones.

SIR W. MARTIN CONWAY has sent to the *Daily Chronicle* the news that on September 9 he reached the top of Yllimani, a peak of the Cordilleras which rises behind the town of La Paz, at a height of 22,500 feet above sea-level. With Sir Martin Conway are the two Swiss guides, Antoine Maquiguez and Louis Pellissier, who last year were in Alaska with the Duke of the Abruzzi, and made the ascent of Mount St. Elias. The party took five days to reach the top of Yllimani from the highest point of cultivation.

MRS. HUBBARD has sent us the following translation of a passage from the "Niva," recording an interesting observation: "The naturalist, Ostrovomov, director of the biological station at Sevastopol, last summer made some excursions along the coast of the Crimea. One morning, the sea being at the time calm and clear and the sky blue, he observed whole clouds of small creatures, like moths, fluttering above the smooth surface of the sea. Ostrovomov, with his son and a boy from the station, observed that each of these small creatures rested for a while on the surface of the water, as though gathering strength; then made a spring and flew high in the air, and plunged again into the sea. They captured some of these, and examined them under the microscope; and what was Ostrovomov's astonishment on discovering that these flying creatures were the soft-shelled crablike *Entomostraca*, belonging to the family *Pontellina mediterranea*."

THE world's record for high kite flight was (says *Science*) broken on August 26 at Mr. Rotch's observatory by Messrs. Clayton and Ferguson, who despatched a tandem of kites into the air until the highest one reached an altitude of 12,124 feet above the sea-level, a height 277 feet greater than any kite had previously reached. Five miles of line, weighing 75 pounds, was let out, while the weight of the kites, recording instruments and secondary line, was 37 pounds, making a total of 112 pounds lifted into the air. The recording instrument was made by Mr. Ferguson and was of aluminum, weighing three pounds, and registering temperature, pressure, humidity and wind velocity. The ascent was begun at 11 o'clock, and the highest point reached at 4.15 p.m. The kites passed through clouds when about a mile above the surface of the earth, but while above the clouds the instruments showed the air to be very dry. At the highest point the temperature had fallen to 38°, and the wind velocity was 32 miles an hour. At the ground at the same time the temperature was 75° and the wind velocity 32 miles. The highest wind velocity recorded was 40 miles an hour at a height of 11,000 feet. The wind on the ground at this time was from the west, while at the highest point reached by the kites it was south-west. The flight was one of a series of high ascents made during the spring and summer, averaging about a mile and a half, while on several occasions a height of over 10,000 feet has been obtained.

THE results of meteorological observations made at Rousdon Observatory, South Devon, under the superintendence of Sir Cuthbert Peek, have been published. Interest in local meteorology is necessarily limited, but there are several sections of the present report which appeal to meteorologists generally. The usual comparison was made of daily forecasts issued by the Meteorological Office for the district in which the Observatory is situated with actual weather experienced. The wind and weather predictions were both correct in 85 per cent. of the forecasts. The forecasts of wind alone were correct in 90 per cent., and 92 per cent. of the weather predictions were fulfilled. The percentage of correct weather forecasts has not been below 92 for the past five years. Sir Cuthbert Peek has made a further comparison of the records of the Robinson cup anemometer and the pressure-tube anemometer. It has been assumed that the factor of the cup anemometer does not depend upon, or vary with, the velocity of the wind. To roughly test this conclusion, a comparison was made of the daily total mileages of wind passing over the Observatory, as recorded by the two instruments, during three periods of about twenty-four days each, when light airs, winds of moderate force, and strong winds, respectively, prevailed. These results show quite clearly the effect of the inertia of the cups in low velocities, the excess of the cup record over that of the pressure-tube being as much as 53 per cent., when the mean hourly velocity is as low as four miles. With a moderate wind of eleven miles per hour, however, the cups yield 3 per cent. less than the pressure-tube; and with a wind of double that velocity the difference is increased to 8 per cent. It is pointed out that these results are based on too few observations to be accepted as final, but they are suggestive, and a fuller comparison on the lines indicated may at some future time be carried out. The factor 2.2 appears, however, to be practically correct for all winds, except when the force is extremely low.

THE two Cantor Lectures delivered before the Society of Arts by Dr. D. Morris, C.M.G., on sources of commercial india-rubber, have been published in a pamphlet form. In his lectures, Dr. Morris confined himself to describing the rubber plants now existing in various parts of the tropics, their geographical distribution, the conditions under which they grow, and the prospects they afford of being able to meet the increasing demand for rubber. At the outset he made a comparison between india-rubber and gutta-percha. It is very well known that india-rubber and gutta-percha are closely allied substances, not only in their origin but also in their chemical composition. They are both obtained from the latex of certain plants, and are composed wholly of carbon and hydrogen. But, as Dr. Morris points out, the similarity ends here. The most conspicuous property of gutta-percha is that of becoming soft and plastic on immersion in hot water, retaining any shape then given to it on cooling, when it becomes hard and rigid. Caoutchouc, on the other hand, does not soften in moderate heat, is impervious to water, alcohol, most acids, and gases, and retains for a long period its original elasticity and strength. Again, gutta-percha is obtained only from large trees belonging to one family of plants, the *Sapotaceae*, confined to one small portion of the world's surface. Caoutchouc, on the other hand, is obtained from numerous families of plants, and these are distributed over almost every part of the tropical regions; they may be low herbaceous plants, shrubby climbers, small trees, or majestic giants of the forest, 150 to 180 feet high. Dr. Morris's lectures deal with these plants with special reference to the rubber industries connected with our Colonial and Indian possessions; they are, therefore, of great interest at the present time.

In the *Bulletin International* of the Imperial Academy of Science of Bohemia, Herr Franz Švec discusses the ciliated
NO. 1508, VOL. 58]

Infusoria of the Unterpočernitzer Teich, on the banks of which a biological station has been established. Sixty-nine species of Infusoria have been observed, of which ten are new; these are *Holophyra atra*, *Enchelys variabilis*, *Lacrymaria phialina*, *Pyradon nucleatus*, *Lionotus lanceolatus*, *Loxophyllum aselli*, *Dileptus elephantinus*, *Zoothamnium limneticum*, *Epistylis rotans*, *Rhabdostyla discostyla*.

THE "Communications from the Physical Laboratory at the University of Leiden," published in English under the direction of Prof. Kamerlingh Onnes, afford a striking instance of the activity of foreign Universities in the matter of research. Part 41 contains a paper by Dr. E. van Everdingen, jun., on the Hall-effect in electrolytes. The author has calculated the amount of the effect in liquids, and has compared the results with those afforded by experiments, but it appears that the observed galvanometric differences of potential in liquids differ considerably (sometimes even in sign) from those which would be caused by the Hall phenomenon. For the present, it therefore appears that we cannot use the phenomenon in electrolytes to obtain a better insight into the nature of the electric current in metals.

THE *Proceedings* of the Royal Society of Queensland (vol. xiii.), just published, comprise several papers of interest. In a useful presidential address, Mr. C. J. Pound shows how the stockowner is indebted to the microscope, and explains that "all those marvellous and brilliant discoveries relating to the origin, nature, prevention, and treatment of bacterial diseases of our domesticated animals have been mainly brought about by the investigations of such brilliant epoch-making men as Pasteur, Koch, and Lister, whose names will ever be associated with the microscope and remain as lasting monuments to the science of preventive medicine."—Mr. Walter E. Roth contributes some notes on social and individual nomenclature among certain north Queensland aborigines, personally studied by him. Mr. Roth points out that the whole question of class-systems, whereby a relationship, such as it is, is established between aborigines living miles and miles apart, yet may be mutually unknown personally, has an important practical bearing which has hitherto been apparently overlooked. In the mind of the real North-west Central Queensland savage, all white men are believed to be similarly related; he looks upon any one European as being the brother, brother-in-law, father, or mother's brother, &c., of any other.—Mr. Thomas P. Lucas gives descriptions of Queensland lepidoptera, and Mr. Rowland Illidge contributes a list of butterflies of the Brisbane district.

A BRIEF statement of the results of an investigation into the distribution and ethnography of leprosy in the Far East is given by Mr. Sydney B. J. Skertchly in the volume of *Proceedings* referred to in the foregoing note. The area embraced in an inquiry carried out by Mr. Skertchly and Dr. J. Cantlie, extends from the Malay Peninsula, through China, the whole of the East Indian Archipelago, Japan, and the Philippines and the islands of the Pacific; and a large amount of most valuable information as to the distribution of the disease has been obtained. As the area investigated contained every variety of surface, it was easy to determine whether physical configuration was a determining cause of leprosy. The conclusion arrived at is that neither physiographical climate nor geological conditions have any influence upon the distribution of leprosy. Contrary to the general statements, leprosy is not rife throughout the length and breadth of China, entire provinces being free from the disease. An examination of the state of affairs in the Pacific leads to the important conclusion that from the Chinese provinces of Kwantung and Fokien, leprosy spreads with diminishing intensity in all directions, and has formed a new focus in Hawaii of unparalleled virulence. Viewing the facts

from an anthropological standpoint, it appears that so far from the black races being the most leprosy, and the yellow the least, over the great area dealt with, the black races are quite free from leprosy, except where, as in Fiji, it has been recently introduced; and the yellow race, the Chinese, is the leper and the distributor of leprosy. In not a single instance are the native races attacked without there being Chinese lepers in the country. In other words, leprosy follows the lines of Chinese emigration, and in the East Indian Archipelago and Oceania is co-terminous and co-existent, in time and area, with the Chinese coolie. Mr. Skerchly believes that the only way to stop the spread of leprosy is to put an end to the coolie traffic from the infected provinces, and this cannot be done except by concerted action of the Governments holding possessions in the Far East.

THE Wilde lecture "On the Physical Basis of Psychical Events," delivered by Prof. Michael Foster before the Manchester Literary and Philosophical Society last March, is printed in *Manchester Memoirs*, vol. xlii. (1898), No. 12.

A COPY of "Bourne's Handy Assurance Manual" (1898), edited by Mr. William Schooling, has been received. The volume shows the position of every assurance office, and should be consulted before taking out a policy in any company. Students of statistics will also find the tables useful.

A NINTH edition of Skerchly's "Geology," revised in accordance with the latest requirements of the Science and Art Department's syllabus, has been prepared by Dr. J. Monckman, and published by Mr. Thomas Murby. A new section dealing with minerals and their microscopic characteristics has been added, but the general appearance of the book and the illustrations are behind the times.

THE additions to the Zoological Society's Gardens during the past week include a Ring-tailed Coati (*Nasua rufa*) from South America, presented by Mr. S. C. Rogers; two Little Armadillos (*Dasypus minutus*) from Patagonia, a Vociferous Sea Eagle (*Haliastur vocifer*), a Chameleon (*Chamaeleo vulgaris*) from Africa, deposited; a Pleasant Antelope (*Tragelaphus gratus*, ♀), bred in Amsterdam, purchased; a Crested Porcupine (*Hystrix cristata*), three Swinhoe's Pheasants (*Euplocamus swinhoi*), three Mandarin Ducks (*Aix galericulata*), bred in the Gardens.

OUR ASTRONOMICAL COLUMN.

THE NEBULA OF ANDROMEDA.—A telegram from the Centralstelle, Kiel, received here on the 20th, announces that Seraphimoff has observed a stellar-like condensation near the centre of the nebula of Andromeda.

This is not the first time that variations near the centre of this nebula have been observed. In 1885 a star of 6.5 mag. appeared suddenly near the centre, giving a continuous spectrum containing probably a few bright lines; in 1886 this had entirely disappeared. Espin thought that the nucleus was variable, and that he could see stars in it; and Young, with a 23-inch refractor, confirmed this. The fine series of photographs taken by Roberts also indicate that the nucleus of the nebula is variable.

An examination of the nebula on the early morning of the 21st, with the 30-inch reflector of the Solar Physics Observatory, South Kensington, gave the idea that the centre of the nucleus seemed more elongated and was more of a stellar nature than usual. The application of the spectroscopic indicated nothing more than a continuous spectrum, although there may have been faint bright lines which could not be seen.

COMETS TEMPEL 1866 and PERRINE-CHOFARDET.—Just after we had gone to press last week we received another telegram, concerning the comet discovered by Pechuele, saying that it was Wolf's comet and not that of Tempel.

Another telegram, dated September 15, informs us that Perrine, on September 13, discovered a comet at 16h. 14.3m. Lick Mean Time, in position of R.A. 9h. 41m. 40s. and Declination + 30° 36'. Two circulars from Kiel (Nos. 11 and 12), which have since reached us, give the elements of the comet's orbit and an ephemeris for the present month, besides telling us that Chofardet made the same discovery independently at Besançon on September 14, 16h. 37m. local time.

Both the elements calculated by Berberich from observations on September 12, 13, 15, and by Perrine and Aitken from observations on September 13, 14 and 15 are very similar, so we will confine ourselves to the former, which are namely:—

$$T = 1898 \text{ October } 19.9565 \text{ Berlin M.T.}$$

$$e = 165 \text{ } 56' 29''$$

$$\Omega = 36 \text{ } 20' 85''$$

$$i = 29 \text{ } 16' 41''$$

$$\log q = 9.57608$$

For the present month the positions of the comet for every two days are as follows:—

1898.		12h. Berlin M.T.		Decl.	Br.
		R.A.	h. m. s.		
Sept.	22	...	10 35 2	...	+25 9'9
"	24	...	10 48 3	...	23 32'4
"	26	...	11 1 11	...	21 46'6
"	28	...	11 14 23	...	+19 52'3

CATALOGUE OF NEBULÆ.—Mr. Lewis Swift publishes in a recent number of *Astr. Nachr.* (No. 3517) a catalogue of nebulae which have been discovered by him during the last three years. All the observations were made at the Lowe Observatory, Echo Mountain, California, the low latitude of this station, namely + 34° 20', enabling him to search further south than when he was situated at Rochester, New York. He says: "I am further south than any observatory in Europe and America north of the equator except the one at Tacubaya, Mexico, yet I find that the southern sky has been pretty thoroughly explored by Sir John Herschel, Dunlop, and others."

The present catalogue contains 243 objects, some of which are very interesting. Thus, Nos. 6 and 27 are described as being very singular. They resemble a fairly bright double star, each component being an exceedingly small nebulous disc "like an imaginary double nebulous Uranus distant about 5" or 6".

No. 56 is described as "a nebulous hair-line of one uniform size from end to end," while No. 91 has one side extending like a brush.

In addition to the above, this keen-eyed observer has discovered no less than four comets, one of which is of short period, and his son has discovered a fifth, also of short period.

UNIVERSITY AND EDUCATIONAL INTELLIGENCE.

THE Calendar of University College, Bristol, for the Session 1898-99 has been published. The College offers excellent opportunities for the study of science, languages, history and literature, and possesses good facilities for giving systematic instruction in the branches of applied science more nearly connected with the arts and manufactures. Medical education is provided by the Faculty of Medicine of the College; and students can complete in Bristol the entire course of study required for the various medical and surgical degrees.

In order to encourage systematic study, a definite course of instruction extending over three years has been established in the Morley Memorial College (for working men and women), Waterloo Bridge Road. In the first and second years all students will follow almost the same course of study, but in the third year they will take up a selected group of literary, mathematical, or scientific subjects. Organised courses of this kind are of far greater educational value than the study of a large number of disconnected subjects.

MAJOR P. G. CRAIGIE's annual report to the Board on Agriculture on the distribution of grants for agricultural education and research in 1897-98, has just been issued as a Parliamentary paper. The total amount distributed during the financial year to each of the fifteen institutions receiving assistance was 7200*l.*, as compared with 7000*l.* in the previous year. The following table shows how this money was expended:—

Institutions aided.	Work.	Grant, 1897-98.
University College of North Wales, Bangor	Collegiate centre	£ 800
University College of North Wales, Bangor	College farm	200
Durham College of Science, New- castle-on-Tyne	Collegiate centre	800
Durham College of Science, New- castle-on-Tyne	College farm	200
University College of Wales, Aber- ystwyth	Collegiate centre	800
Reading College	Collegiate centre	800
Yorkshire College, Leeds	Collegiate centre	600
University College, Nottingham	Collegiate centre	600
South-Eastern Agricultural College, Wye	Collegiate centre	600
Cambridge and Counties Agricul- tural Education Committee	Collegiate centre	500
Eastern Counties Dairy Institute, Ipswich	Dairy instruction	300
British Dairy Institute, Reading	Dairy instruction	300
Royal Botanic Garden, Edinburgh	Class for foresters and gardeners	150
Bath and West and Southern Coun- ties Society	Field experiments	50
Bath and West and Southern Coun- ties Society	Cider experiments	50
Bath and West and Southern Coun- ties Society	Cheddar cheese research	200
Highland and Agricultural Society Agricultural Research Association, Aberdeen	Agricultural experiments	100
Stewartry of Kircudbright Dairy Association	Cheese discoloration inquiry	50

The grants to the collegiate centres in England and Wales are of a general character, intended to assist and improve the local provision made for instruction in the higher forms of agricultural education. The thirty-two separate counties are thus provided with an efficient and economical means of systematising their local instruction, and of supervising demonstration plots and agricultural experiments by securing scientific advice and the assistance of qualified lecturers drawn from the collegiate educational staffs. The Durham College of Science and the University College of North Wales have been granted special assistance in consideration of their having taken farms for practical work and field experiments.

SOCIETIES AND ACADEMIES.

PARIS.

Academy of Sciences, September 12.—M. Faye in the chair.—Meadow land in warm dry summers, by M. Ad. Chatin. A list of those species of plants which have been found to be the most capable of resisting a hot, dry summer.—Observation of an aurora borealis, by M. H. Deslandres. An aurora was observed at Meudon on September 9 about 9 p.m., and its general direction was very nearly that of the magnetic meridian, the rays having a greenish colour.—On the crystallisation of the anhydrous sulphides of calcium and strontium, by M. Mourlot. The crystallised sulphides of these metals can be prepared in two ways, either by heating a mixture of the corresponding sulphate with carbon, or by simply fusing the anhydrous sulphide obtained by the method of M. Sabatier, the temperature employed being that of the electric furnace with a current of 1000 amperes at 60 volts. The crystallised sulphides thus produced are more stable than the corresponding amorphous salts, and are attacked with difficulty by reagents; carbon at a very high temperature converts them into carbides. Both crystallise in the cubic system, and are without action upon polarised light.—On a double carbide of iron and tungsten, by M. Percy Williams. This compound, the existence of which was indicated in an earlier paper, is prepared by heating a mixture of tungstic acid, iron and coke, in the electric furnace with a current of 900 amperes at 45 volts. The ingot formed in the reaction contains the carbide of tungsten WC, probably W_2C , and the double carbide $3W_2C \cdot 2Fe_3C$.—On the commercial extraction of thorium, by MM. Wyronhoff and A. Verneuil. The mineral is worked up by one of the usual methods as far as the production of the oxalates, these precipitated by sodium carbonate and hydroxide, and the washed precipitate dissolved in hydrochloric acid. This liquid is treated with small portions of barium peroxide, until hydrogen peroxide no longer gives a precipitate. The deposit, which is

of a reddish orange colour owing to the presence of cerium, contains the whole of the thoria, with about 20 to 30 per cent. of impurities. Further treatment with hydrogen peroxide after a similar set of operations readily gives a very pure thoria. The method has been applied on the large scale, starting with five tons of monazite, with good results.—On the composition of the humic constituents of the soil, by M. G. André.—On the transformation of luminous variations into mobile relief, by M. Dussaud.—On a new coccus, by M. Louis Leger. The new species is found in the digestive tube of *Lithobius hexodus*, and belongs to the genus *Echinospira*. Its microgametes are furnished with vibratile cilia; the name *E. ventricosa* is suggested.—Influence of light on the form and structure of the branches of the wild grape and ground ivy, by M. Maige. Comparative cultures placed in light of decreasing intensities showed that both from the morphological and anatomical points of view, a feeble light increases the adaptive powers of climbing plants, diffused light favouring the conversion of a flower-bearing bud into a tendril. Direct sunlight produces the opposite effect.—On the adherence of the cupric solutions used for curing the cryptogamous diseases of the vine, by MM. Guillon and Gourand.

BOOKS RECEIVED.

BOOKS.—The Unconscious Mind: Dr. A. T. Schofield (Hodder).—U.S. Department of Agriculture: Report of the Chief of the Weather Bureau, 1896-97 (Washington).—Bird Studies: W. E. D. Scott (Putnam).—Coffee and India-rubber Culture in Mexico: M. Romero (Putnam).—The Sphere of Science: Prof. F. S. Hoffman (Putnam).—A Text-Book of General Astronomy: Prof. C. A. Young, new edition (Arnold).—A Pocket Dictionary of Hygiene: C. T. Kingzett and D. Homfray (Baillière).—University College, Bristol, Calendar, 1898-99 (Bristol).—A Memoir of T. Sterry Hunt: J. Douglas (Philadelphia).—Infinitesimal Analysis: Prof. W. B. Smith, Vol. 1 (Macmillan).—Die Photometrie der Gestirne: Prof. J. Müller (Leipzig, Engelmann).—Die Photographie der Gestirne: Prof. J. Scheiner (Leipzig, Engelmann).—Atlas ditto (Leipzig, Engelmann).—Untersuchungen zur Physiologie der Pflanzlichen Organisation: Prof. G. Berthold, Erster Teil (Leipzig, Engelmann).—A Text-Book of Geodetic Astronomy: I. T. Hayford (Chapman).


CONTENTS.

PAGE

The Fauna and Flora of the Pamir. By W. T. B.	493
Sociological Science. By A. E. Taylor	494
Our Book Shelf:—	
Strasburger, Noll, Schenck, Schimper: "A Text-Book of Botany"	494
Dumont: "Automobiles sur Rails"	495
Lecornu: "Régularisation du Mouvement dans les Machines"	495
Kingzett and Homfray: "A Pocket Dictionary of Hygiene"	495
Campion: "The Secret of the Poles"	495
Kerr: "Wireless Telegraphy, popularly explained"	495
Letters to the Editor:—	
Chance or Vitalism.—Prof. Karl Pearson, F.R.S.	495
The Moon's Course.—Sir Samuel Wilks, Bart., F.R.S.	496
The Aurora of September 9.—Hon. Rollo Russell	496
A White Sea.—James W. Barrett	496
Deep-Sea Dredging, and the Phosphorescence of Living Creatures, at Great Sea Depths.—E. L. J. Ridsdale	497
The Injection of Cocaine as a Remedy for Stings.—M.D., Oxon	497
The Geography of the United States. (Illustrated.) By Dr. Hugh Robert Mill	497
The British Association	498
Section D.—Zoology. (With Diagrams.)—Opening Address by Prof. W. F. R. Weldon, F.R.S., President of the Section	499
Section G.—Mechanical Science.—Opening Address by Sir John Wolfe Barry, K.C.B., F.R.S., President of the Section	506
International Sea Fisheries Congress at Dieppe	511
Notes	512
Our Astronomical Column:—	
The Nebula of Andromeda	515
Comets Tempel 1866 and Perrine-Chofardet	515
Catalogue of Nebulae	515
University and Educational Intelligence	515
Societies and Academies	516
Books Received	516

W. G. PYE,
Physical & Physiological Instrument Maker,
30 ST. ANDREW'S STREET, CAMBRIDGE.

STUDENTS'
BRIDGE
Aggregate
Guaranteed
Platinoid



WHEATSTONE
BOX.
12,220 ohms
to 1 in 500.
Wire.

Price £9 10s.

IMPROVED REFLECTING GALVANOMETER,
for use with above, any resistance to 2000 ohms, £3.
Galvanometer Shunts, Galvanometer Lamps and Scales,
Keys, Single Coils, Batteries, &c.

Reliable Apparatus at low prices for teaching Mechanics,
Hydrostatics, Heat, Optics, Magnetism and Electricity, as
supplied to many of the leading Technical Institutes and
Organised Science Schools.

Lists Free on application. Physiologic List in preparation.



FREDK. JACKSON & Co.
(Late MOTTESHEAD & CO.),

14 CROSS STREET, MANCHESTER

Goods Entrance: 10 Half-Moon Street,

LABORATORY FURNISHERS,
Importers, Manufacturers, and Dealers in
**CHEMICAL AND PHYSICAL
APPARATUS**

Of every Description.
Fine Chemicals, Volumetric Solutions,
Plain and Stoppered Bottles,
AND EVERY LABORATORY REQUIREMENT.
Illustrated Catalogue of Apparatus, with Price List
of Chemicals, free on application.

Telegraphic Address—"APPARATUS, MANCHESTER."
Telephone Number—2038.

BREWSTER, SMITH, & CO.,
6 CROSS STREET, FINSBURY PAVEMENT, E.C.

MANUFACTURERS AND IMPORTERS OF
CHEMICAL AND SCIENTIFIC APPARATUS

AND
PURE CHEMICALS.

Apparatus for School and Trade purposes.

Finest Bohemian and Jena Hard Glass, also Cheap German
Glass Goods, graduated and verified Burettes, Pipettes, and
c.c. Measures, Balances, Metric Weights, either separately or
in Sets.

Our New Test Tube Stands (Patent applied for) will hold any size Test
Tubes, from $\frac{1}{8}$ to $1\frac{1}{2}$ inch diameter. A great advantage over the ordinary
Stands in use, which only have $\frac{1}{2}$ -inch holes.

Price Lists ready about September 20.

**RÖNTGEN TUBES WITH VACUUM
REGULATOR.**

(PROF. ZEHNDER'S PATENT.)

A GREAT SUCCESS!

SOLE AGENTS:


WILHELM & CO.,

11 & 12 WESTMORLAND BUILDINGS, ALDERSGATE STREET,
LONDON, E.C.

C. E. MÜLLER & CO.,

MANUFACTURERS OF

CHEMICAL AND PHYSICAL APPARATUS,
X-Ray Tubes, Argon and Helium Spectrum Tubes,
Special Apparatus for Research, Chemical
Thermometers, &c., &c.



**NEW DOUBLE
SURFACE
CONDENSER,**

designed to take the place
of the ordinary Liebig
Condenser. Whilst being
more efficient, it is *only*
one-fifth the size.

**SETS OF APPARATUS
FOR
STUDENTS' USE.**

Price Lists and full particulars of above on application.

148, HIGH HOLBORN, LONDON, W.C.

**THOMAS'S
PLATES AND FILMS,**

ORDINARY or ISOCHROMATIC,
IN THREE RAPIDITIES.

WHITE LABELS.

MEDIUM ORDINARY.

EXTRA RAPID do.

"A1" do.

YELLOW LABELS.

MEDIUM ISOCHROMATIC.

EXTRA RAPID do.

"A1" do.

The Extra Rapid are three times and "A1" six times faster than the Medium.

PLATES $\frac{1}{4}$ $\frac{1}{2}$ $\frac{3}{4}$ $\frac{4}{3}$

FILMS $\frac{1}{6}$ $\frac{3}{8}$ $\frac{1}{2}$ $\frac{2}{3}$

R. W. THOMAS & Co., Ltd.,

PALL MALL FACTORY,

THORNTON HEATH.

BAIRD & TATLOCK,

Manufacturers of Chemical and Physical
Apparatus, Laboratory Benches, Fume
Closets, &c.

Dealers and Importers of Pure
Chemicals, &c.

Contractors to Her Majesty's Government.

14 CROSS STREET, HATTON GARDEN,
LONDON, E.C.

CATALOGUES ON APPLICATION.

MACMILLAN & CO.'S BOOKS

FOR

STUDENTS OF ZOOLOGY.

A TEXT-BOOK OF ZOOLOGY. By T. JEFFERY PARKER, D.Sc., F.R.S., Professor of Biology in the University of Otago, Dunedin, N.Z., and WILLIAM A. HASWELL, M.A., D.Sc., F.R.S., Professor of Biology in the University of Sydney, N.S.W. In Two Vols. With Illustrations. Medium 8vo, 36s. net.

LITERATURE.—"The scheme of classification is on the whole so good that even those who find fault with matters of detail will admit its general excellence. The sections on Distribution—geographical, bathymetrical, and geological—and on The Philosophy of Zoology have been written with extreme care, the latter especially being full of pregnant remarks and suggestions. A useful chapter on the History of Zoology is appended, from which a true idea of the perspective of things can be gathered. Both volumes are copiously indexed, and reference to any heading is thus made easy."

A TEXT-BOOK OF ENTOMOLOGY, including the Anatomy, Physiology, Embryology and Metamorphoses of Insects, for use in Agricultural and Technical Schools and Colleges, as well as by the working Entomologist. By ALPHEUS S. PACKARD, M.D., Ph.D., Professor of Zoology and Geology, Brown University. Medium 8vo, 18s. net.

NATURE.—"Dr. Packard's book will be of the greatest service to students of insect-anatomy, and almost indispensable to future writers on the subject. . . . Should find a place in every library which includes comparative anatomy, and should be the constant companion of all who occupy themselves with the structure and life-histories of insects."

ESSAYS ON MUSEUMS AND OTHER SUBJECTS CONNECTED WITH NATURAL HISTORY. By Sir WILLIAM HENRY FLOWER, K.C.B., D.C.L., D.Sc., LL.D., Ph.D., F.R.S., F.R.C.S., P.Z.S., Correspondent of the Institute of France, &c. Demy 8vo, 12s. net.

ATHENÆUM.—"Many of the essays possess permanent value, and all were well worthy of being printed afresh."

A TEXT-BOOK of COMPARATIVE ANATOMY. By Dr. ARNOLD LANG, Professor of Zoology in the University of Zürich. With Preface to the English Translation by Dr. ERNST HAECKEL. Translated by HENRY M. BERNARD, M.A. Cantab., and MATILDA BERNARD. Vols. I. and II. (completing the Invertebrates). 8vo. 17s. net each.

LESSONS in ELEMENTARY ANATOMY. By ST. GEORGE MIVART, F.R.S., Author of "The Genesis of Species." Fcap. 8vo. 6s. 6d.

ELEMENTS of the COMPARATIVE ANATOMY OF VERTEBRATES. Adapted from the German of ROBERT WIEDERSHEIM, Professor of Anatomy, and Director of the Institute of Human and Comparative Anatomy in the University of Freiburg, in Baden. By W. NEWTON PARKER, Professor of Biology in the University College of South Wales and Monmouthshire. With additions by the Author and Translator. 270 Woodcuts. Medium 8vo. Second Edition. 12s. 6d. net.

The STRUCTURE of MAN: An INDEX to his PAST HISTORY. By ROBERT WIEDERSHEIM. Translated by H. and M. BERNARD. The Translation, Edited and Annotated, and a Preface written by G. B. HOWES, F.R.S., Professor of Zoology, Royal College of Science, London. With 105 Figures in the Text. 8vo. 8s. net.

A COURSE of INSTRUCTION in ZOOTOMY. Vertebrata. By T. JEFFERY PARKER, F.R.S., Professor of Biology in the University of Otago, New Zealand. With Illustrations. Crown 8vo. 8s. 6d.

An INTRODUCTION to the OSTEOLOGY of the MAMMALIA. By Sir WILLIAM HENRY FLOWER, F.R.S., F.R.C.S., Director of the Natural History Department of the British Museum. Illustrated. Third Edition. Revised with the assistance of HANS GADOW, Ph.D., Lecturer on the Advanced Morphology of Vertebrates in the University of Cambridge. Crown 8vo. 10s. 6d.

A TREATISE on COMPARATIVE EMBRYOLOGY. By the late Prof. F. M. BALFOUR, M.A., F.R.S. In two Vols. Second Edition. Medium 8vo. Vol. I. 18s. Vol. II. 21s.

The ELEMENTS of EMBRYOLOGY. By Prof. MICHAEL FOSTER, M.D., F.R.S., and the late Prof. F. M. BALFOUR, F.R.S. Second Edition, revised. Edited by A. SEDGWICK, M.A., and W. HEAPE, M.A. Illustrated. Crown 8vo. 10s. 6d.

STRUCTURE AND LIFE of BIRDS. By F. W. HEADLEY, Assistant Master at Haileybury College. Illustrated. Crown 8vo. 7s. 6d.

The NATURAL HISTORY of the MARKETABLE MARINE FISHES of the BRITISH ISLANDS. By J. T. CUNNINGHAM, M.A. With Preface by E. RAY LANKESTER, F.R.S. Illustrated. Medium 8vo. 7s. 6d. net.

NATURAL HISTORY of AQUATIC INSECTS. By L. C. MIALL, F.R.S., Professor of Biology in the Yorkshire College, Leeds. Illustrated. Crown 8vo. 6s.

THE ORIGIN AND METAMORPHOSES of INSECTS. By Sir JOHN LUBBOCK, F.R.S. With Illustrations. 3s. 6d. [*Nature Series.*]

MONOGRAPH of the BRITISH CICADÆ or TETTIGIIDE (Froghoppers and Grass-flies). By GEORGE BOWDLER BUCKTON, F.R.S., Illustrated by more than 400 Coloured Drawings. Two Vols. Demy 8vo. 33s. 6d. each, net. Also in Eight Parts. 8s. each, net.

A HANDBOOK of BRITISH LEPIDOPTERA. By EDWARD MEYRICK, B.A., F.Z.S., F.E.S., Assistant Master at Marlborough College. Extra Crown 8vo. 10s. 6d. net.

MACMILLAN AND CO., LIMITED, LONDON.

MACMILLAN'S ELEMENTARY SCIENCE CLASS BOOKS.

Adapted to the Elementary Stages of the South Kensington Syllabuses.

JUST PUBLISHED. Globe 8vo, 2s. 6d.

GEOLOGY FOR BEGINNERS. By W. W. WATTS, M.A., F.G.S., formerly Lecturer to the Cambridge University Extension, and Member of Her Majesty's Geological Survey, &c. With 310 Illustrations.

GUARDIAN.—"Since Geology has begun to take its place in school curricula a work has been needed adapted to junior classes, and one which lends itself to 'getting up' for examinations. The book before us seems well designed to supply this need. . . . Abundant illustrations and a copious index add to the value of a book which is evidently the work of one who knows the ground."

PRACTICAL PLANE and SOLID GEOMETRY. With an Introduction to Graphic Statics. By J. HARRISON, M.Inst.M.E., &c., Whitworth Scholar; and G. A. BAXANDALL. 2s. 6d.

ARITHMETIC FOR SCHOOLS. By S. L. LONEY, M.A. With or without Answers. 4s. 6d. each.

ARITHMETIC FOR SCHOOLS. By Rev. J. B. LOCK, M.A. New Edition. 4s. 6d. Key, 10s. 6d.

A TEXT-BOOK OF EUCLID'S ELEMENTS. By H. S. HALL, M.A., and F. H. STEVENS, M.A. Book I., (1s. Books I. to IV., 3s. Complete, 4s. 6d. Key to Books I.-IV., 6s. 6d. Key to V., VI., and XI., 3s. 6d. Complete, 8s. 6d.

ALGEBRA FOR BEGINNERS. By H. S. HALL, M.A., and S. R. KNIGHT, B.A. 2s. With Answers, 2s. 6d.

[This book deals with Algebra as far as Quadratic Equations.]

ELEMENTARY ALGEBRA. By H. S. HALL, M.A., and S. R. KNIGHT, B.A. 3s. 6d. With Answers. 4s. 6d. Key, 8s. 6d.

[This book covers the Second and Third Stages.]

ALGEBRA FOR BEGINNERS. By I. TOD-HUNTER. New Edition, thoroughly Revised by S. L. LONEY, M.A. Globe 8vo. 3s. 6d. With Answers, 4s. 6d. Key, 8s. 6d. net.

MECHANICS FOR BEGINNERS. By Rev. J. B. LOCK. Part I. Mechanics of Solids, 2s. 6d.

MECHANICS FOR BEGINNERS. By W. GAL-LATLY, M.A. 2s. 6d.

HYDROSTATICS FOR BEGINNERS. By F. W. SANDERSON, M.A. 2s. 6d.

ELEMENTARY LESSONS IN HEAT, LIGHT, AND SOUND. By Prof. D. E. JONES, B.Sc. 2s. 6d.

ELECTRICITY AND MAGNETISM FOR BEGINNERS. By F. W. SANDERSON. 2s. 6d.

AN INTRODUCTION TO THE STUDY OF CHEMISTRY. By W. H. PERKIN, JUN., Ph.D., F.R.S., Professor of Chemistry in Owens College, and B. LEAN, D.Sc. 2s. 6d.

INORGANIC CHEMISTRY FOR BEGINNERS. By Sir HENRY ROSCOE, F.R.S., and JOSEPH LUNT, B.Sc. New Edition, 1895. 2s. 6d.

PRACTICAL INORGANIC CHEMISTRY. By G. S. TURPIN, M.A., D.Sc. 2s. 6d.

A JUNIOR COURSE OF PRACTICAL CHEMISTRY. By F. JONES, F.C.S. With Preface by Sir H. E. ROSCOE, F.R.S. Eighth Edition. 2s. 6d.

ORGANIC CHEMISTRY FOR BEGINNERS. By G. S. TURPIN, M.A., D.Sc. 2s. 6d.

PHYSIOLOGY FOR BEGINNERS. By Prof. M. FOSTER and Dr. SHORE. 2s. 6d.

PHYSIOGRAPHY FOR BEGINNERS. By A. T. SIMMONS, B.Sc. (Lond.). 2s. 6d.

HYGIENE FOR BEGINNERS. By E. S. REYNOLDS, M.D. 2s. 6d.

MACMILLAN AND CO., LIMITED, LONDON.

NOTICE.—Advertisements and business letters for NATURE should be addressed to the Publishers; Editorial communications to the Editor. The telegraphic address of NATURE is "PHUSIS," LONDON.

SUBSCRIPTIONS TO "NATURE."

	£	s.	d.
Yearly	1	8	0
Half-yearly	0	14	6
Quarterly	0	7	6
TO ALL PLACES ABROAD:—			
Yearly	1	10	6
Half-yearly	0	15	6
Quarterly	0	8	0

CHARGES FOR ADVERTISEMENTS.

	£	s.	d.
*Three Lines in Column	0	2	6
Per Line after	0	0	9
One Eighth Page, or Quarter Column	0	18	6
Quarter Page, or Half a Column	1	15	0
Half a Page, or a Column	3	5	0
Whole Page	6	6	0

* The first line being in heavy type is charged for as Two Lines.

Cheques and Money Orders payable to MACMILLAN & CO., Limited.

OFFICE: ST. MARTIN'S STREET, LONDON, W.C.

SECOND-HAND (By Leading Makers).

STUDENT'S MICROSCOPES.
WATSON EDINBURGH STUDENT'S, £8 10s, £10, & £12 10s.
BECK'S No. 1 INTERNATIONAL STAND, £30.
BECK'S No. 2 INTERNATIONAL, complete with fittings, £25.
ROSS'S LARGEST MODEL STAND, cost £100, complete, £45.
ASTRONOMICAL TELESCOPES, £2 10s. to £120, in stock.
CLARKSON & CO., 28 Bartlett's Buildings, Holborn Circus, E.C.

COSSOR X-RAY TUBES

With Osmium and Osmium - Iridium Anodes as supplied to J. Mackenzie Davidson, Esq.

ACTUAL MAKER,
67 FARRINGTON
ROAD, E.C.

MERCURY PUMPS.

EXPERIMENTAL AND SCIENTIFIC GLASS-BLOWING.

OPTICAL & SCIENTIFIC INSTRUMENTS.


Spectrometers, Spectroscopes, Goniometers, Cathetometers, Optical Benches, &c., &c. Instruments for special purposes constructed to Clients' own designs. *Price List on application.*

W. WILSON (formerly Foreman at Messrs. ELLIOTT BROS.),
56 Crogsland Road, Chalk Farm, London, N.W.

CONTRACTOR TO H. M. GOVERNMENT.

TO SCIENCE LECTURERS. HUGHES' MOTO-PHOTOSCOPE FOR LIVING PICTURES. THE MOST PERFECT.

No SHUTTER, therefore no Flickering. Superb Mechanism.
The Moto-Photo Camera for taking the Pictures. Illustrated Lists, ad. See Mr. HUGHES' PATENT COMBINATION OPTICAL LANTERN, &c. Miniature Triple Lantern constructed for B. J. MALDEN, Esq.; great success. New Oxyhydrogen Microscope. Science Lanterns for Class Demonstration. Magnificent Results. Docwra Triple, Prize Medal, Highest Award. Supplied to the Royal Polytechnic Institution, Dr. H. GRAY, TAN GUINNESS, Madame ADRIANA PATTI, &c., &c. Patent Pampheng's Science Lanterns. The Universal Lantern 4-inch Condensers, 4-wick Lamp. Portrait Combination front Lenses, 18s. 6d., Marvellous value. Science Lecture Sets. Novelties. The Lantern Kaleidoscope. Cheapest Lantern Outfits in the World. Grandly Illustrated Catalogue, over 180 choice Engravings, 6d.; Postage, 3d. List of 300 Lecture Sets, Science Subjects, Views, &c., 6d.; Postage, ad.. Pamphlets Free.—W. C. HUGHES, SPECIALIST, Brewster House, 82 Mortimer Road, Kingsland, N.



PERKEN, SON & RAYMENT, 99 HATTON GARDEN, HOLBORN CIRCUS, LONDON.

LARGE MODEL ILLUSTRATING HERTZIAN WAVE MOTION.

As designed by Prof. SILVANUS P. THOMPSON.

Beautifully illustrating the propagation of the Wave, the progress of which may be watched, from the Oscillator until it reaches the Resonator and causes a responsive vibration.

Apparatus for Hertz Experiments. Radiator and Coherer to Illustrate WIRELESS TELEGRAPHY. Syntonic Leyden Jars with Air-gap Tuning Slide, &c.

HARVEY & PEAK

(BY APPOINTMENT TO THE ROYAL INSTITUTION OF GREAT BRITAIN.
GOVERNMENT CONTRACTORS, &c.,

56 CHARING CROSS ROAD, LONDON, W.C.

NATURAL SCIENCE.

A MONTHLY REVIEW OF SCIENTIFIC PROGRESS.

Contents for OCTOBER 1898:

NOTES AND COMMENTS:—Wanted, an Editor—Biological Work at Bristol—Bread out of Air—Mysteries of Matter—Stereochimistry and Vitalism—Need of Numerical Investigations in Biology—Morphology—Form and Function—Rind Fungus and Sugar Cane—Agriculture in U.S.—Recent Work on the Foraminifera—The Periodical Cicada—Larva of Pelophila—Flat Fish of South Africa—Ichthyosaurus—A New Dinosaur—The Exhibition of Extinct Vertebrata—Cretaceous Rocks in West Greenland—Tertiary and Later Geology of West Greenland.
THE SPECIES, THE SEX, AND THE INDIVIDUAL. (Part II.) By J. T. CUNNINGHAM, M.A., F.Z.S.
BEES AND THE DEVELOPMENT OF FLOWERS. By F. W. HEADLEY, F.Z.S.
THE ESKERS OF IRELAND. (Part II.) By THOMAS FITZPATRICK, LL.D.
THE GREY MULLET FISHERY IN JAPAN. (Illustrated.) By K. KISHINOUE.
THE ZOOLOGICAL CONGRESS.
JAMES HALL.
ANIMAL INTELLIGENCE. By Prof. C. LLOYD MORGAN, F.G.S.
SOME NEW BOOKS:—"Fossil Plants," by A. C. SEWARD; "Bau und Leben unserer Waldbäume," by M. BÜSGEN; "Anthropology of Peru"; "Faune de France," by A. ACOQUE; "Bibliography of Mexican Geology," by R. AGUILAR; "Bibliography of Westralian Geology"; "Hereford Earthquake," by C. DAVISON; "Scraps from Serials, &c." OBITUARIES.—N. A. POMEL—H. CROSSE—F. BERNARD—G. E. GRIMES—Johan Lange, &c.
NEWS.

London: J. M. DENT & CO., 29 & 30 Bedford Street, W.C.

Price One Shilling Net.

Sent Post-free for Annual Subscription of Thirteen Shillings.

THE THORNTON-PICKARD

CAMERAS AND SHUTTERS.

"AMBER" AND "RUBY" CAMERAS FROM £2 13s. 6d.
TIME AND INSTANTANEOUS SHUTTERS FROM 12s. 6d.
Illustrated Catalogue Free.

THE THORNTON-PICKARD MANUFACTURING CO., LTD., ALTRINCHAM.

"OPTIMUS" INDUCTION COILS

(Best London made)

FOR RÖNTGEN X-RAY PHOTOGRAPHY.

Inch Spark	2	3	4	5	6	10	12
Price	£8	£10	£12	£14	£16	£28	£39

Larger Coils to Order.

RADIANT VACUUM **FOCUS TUBES.** RESULTS GUARANTEED. 25/-
ACCUMULATORS AND PRIMARY BATTERIES.

PERKEN, SON & RAYMENT, 99 HATTON GARDEN, HOLBORN CIRCUS, LONDON.

k at
and
ogy
e in
a—
few
cks
l.
II.)

W.
ck,
By

und
":
can
":
es—

.C.

60.
60.

AM.

S

5/-
.

N.

be

SS

N.